

## **Analytical approximation solution of pseudo-parabolic Fractional Equation using a modified doublelaplace decomposition method**

**SOAMPA Bangan<sup>1</sup>, DJIBIBE Moussa Zakari<sup>2</sup> and TCHARIE Kokou<sup>3</sup>**

### **Abstract**

In this paper, the modification of double Laplace decomposition method is proposed for the analytical approximation solution of pseudo-parabolic fractionalequationwithinitialconditions. Someexamplesare given to support the validity and applicability of the presented method. In addition, we check the convergence of double Laplace method applied to our problem.

**Keywords:** Double Laplace transform, inverse double Laplace transform, singular pseudo-parabolic fractional equation, single Laplace transform, decompositionmethods.

---

<sup>1</sup>Mathematics Department, University of Lomé.

<sup>2</sup>Mathematics Department, University of Lomé.

<sup>3</sup>Mathematics Department, University of Lomé.

## 1 Introduction

Fractional differential equations can describe many phenomena in various fields of engineering and scientific disciplines such as control theory, physics, chemistry, biology, economic, mechanics and electromagnetic.

In theoretical physics, it is usually very important to seek and construct explicit solutions of linear and nonlinear partial differential equations (PDEs). Therefore, the solution helps the researchers to understand the physical phenomena. The parabolic equation occurs in several areas of applied mathematics, such as heat conduction, the phenomena of turbulence and flow through a shock wave traveling in a viscous fluid such as the modeling of dynamics. In recent years, several studies for the linear and nonlinear initial value of fractional problems arise in the literature. The pseudo-parabolic equation models a variety of physical processes. The one dimensional pseudo-parabolic equation was derived in [12]. In general, some of the nonlinear models of real-life problems are still very difficult to solve either theoretically or numerically. Recently, many authors have proposed analytical solution to one dimensional system of parabolic equation ( Burgers equation) eg.[13, 14] using a domain decomposition method. In [5] the author used Laplace transform and homotopy perturbation method to obtain approximate solutions of homogeneous and inhomogeneous coupled Berger's equation. The author in [1] used a modified double Laplace decomposition method to solve coupled pseudo-parabolic equations. Our aim in this work, is to use modified double Laplace decomposition for solving a singular one dimensional pseudo-parabolic fractional equation.

## 2 Preliminary Notes

We will now recall the following definitions which are given by [1, 15]

**Definition 2.1.** *The double Laplace transform of the functions  $u(x, t)$  and  $f(x, t)$  are defined as*

$$L_x L_t [u(x, t)] = U(p, s) = \int_0^\infty \int_0^\infty e^{-px} e^{-st} u(x, t) dt dx, \quad (1)$$

$$L_x L_t [f(x, t)] = F(p, s) = \int_0^\infty \int_0^\infty e^{-px} e^{-st} f(x, t) dt dx, \quad (2)$$

where  $x, t > 0$  and  $p, s$  two complex values.

**Definition 2.2.** The Caputo derivative of the function  $u(x, t)$  is defined as follow:

$$\mathbb{D}_t^\alpha v(x, t) = \frac{1}{\Gamma(1 - \alpha)} \int_0^t \frac{\partial v(x, \tau)}{\partial \tau} \times \frac{1}{(t - \tau)^\alpha} d\tau. \quad (3)$$

**Definition 2.3.** The single Laplace transform of Caputo derivative of  $\alpha$  order ( $0 < \alpha < 1$ ) of the function  $f(t)$  is defined as

$$L[\mathbb{D}_t^\alpha f(t)](s) = s^\alpha F(s) - s^{\alpha-1} f(0) \quad (4)$$

where  $F(S)$  is Laplace transforme of  $f(t)$ .

Then, the double Laplace transform of the Caputo derivative of the function  $u(x, t)$  with respect to  $x$  and  $t$  is given by

$$\begin{aligned} L_x L_t [\mathbb{D}_t^\alpha u(x, t)] &= \int_0^\infty \int_0^\infty e^{-px} e^{-st} \mathbb{D}_t^\alpha u(x, t) dt dx \\ &= \int_0^\infty e^{-px} \left[ \int_0^\infty e^{-st} \mathbb{D}_t^\alpha u(x, t) dt \right] dx \\ &= \int_0^\infty e^{-px} \left[ s^\alpha \int_0^\infty e^{-st} u(x, t) dt - s^{\alpha-1} u(x, 0) \right] dx \\ &= s^\alpha U(p, s) - s^{\alpha-1} U(p, 0). \end{aligned} \quad (5)$$

The following basic lemma of the double Laplace transform is given and used in this paper.

**Lemma 2.4.** Double Laplace transform of the non constant coefficient Caputo derivative  $x\mathbb{D}_t^\alpha u(x, t)$  and the function  $xf(x, t)$  are given by

$$\begin{aligned} L_x L_t [x\mathbb{D}_t^\alpha u(x, t)] &= -\frac{d}{dp} [L_x L_t [x\mathbb{D}_t^\alpha u(x, t)]] \\ &= -\frac{d}{dp} [s^\alpha U(p, s) - s^{\alpha-1} U(p, 0)] \end{aligned} \quad (6)$$

$$\begin{aligned}
L_x L_t [x f(x, t)] &= -\frac{d}{dp} [L_x L_t [f(x, t)]] \\
&= -\frac{dF(p, s)}{dp}
\end{aligned} \tag{7}$$

One can prove this lemma by the definition of double Laplace transform in Eq.(2) and Eq.(6).

### 3 Main Results

These are the main results of the paper.

#### 3.1 One Dimensional Pseudo-Parabolic Fractional Equation

In this section we will use modified double Laplace decomposition method to solve singular one dimensional pseudo-parabolic fractional equation. We consider singular one dimensional pseudo- parabolic fractional equation with initial conditions in the form

$$\mathbb{D}_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = f(x, t), \tag{8}$$

subject to Cauchy condition

$$u(x, 0) = g(x), \quad x \geq 0, \quad t > 0, \tag{9}$$

where, the linear term,  $\frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right)$  is called Bessels operator and  $f(x, t)$ ,  $g(x)$  are known functions. In order to obtain the solution of Eq. (9), we apply the following steps.

**Step 1** Multiplying both sides of Eq.(9) by  $x$ .

**Step 2** Using Lemma 2.4 and definition of the double Laplace transform of Caputo derivative for equations in Step 1 and single Laplace transform for initial condition, we get

$$\frac{dU(p, s)}{dp} = \frac{1}{s} \frac{dG(p)}{dp} + \frac{1}{s^\alpha} \frac{dF(p, s)}{dp} - \frac{1}{s^\alpha} L_x L_t \left[ \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) + \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) \right]. \tag{10}$$

where  $G(p), U(p, s), F(p, s)$  are Laplace transform of the functions  $g(x), u(x, t)$  and  $f(x, t)$  respectively.

**Step 3** Applying the integral for both sides of Eq.(10), from 0 to  $p$  with respect to  $p$ , we have

$$U(p, s) = \frac{G(p)}{s} + \frac{1}{s^\alpha} F(p, s) - \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) + \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) \right] dp. \quad (11)$$

**Step 4** Using double Laplace Adomain decomposition methods to define the solution of the singular one dimensional pseudo-parabolic fractional equation as

$$u(x, t) = \sum_{n=0}^{\infty} u_n(x, t). \quad (12)$$

**Step 5** By applying inverse double Laplace transform on both sides of Eq. (11) and use Eq.(12), we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} u_n(x, t) &= g(x) + L_p^{-1} L_s^{-1} \left[ \frac{1}{s^\alpha} F(p, s) \right] \\ &- L_p^{-1} L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial}{\partial x} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right] \\ &- L_p^{-1} L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right]. \end{aligned} \quad (13)$$

In particular, we have

$$u_0 = g(x) + L_p^{-1} L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p dF(p, s) \right] \quad (14)$$

and the rest terms can be written as follows

$$\begin{aligned}
u_{n+1} = & -L_p^{-1}L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial}{\partial x} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right] \\
& - L_p^{-1}L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right], \tag{15}
\end{aligned}$$

where  $L_p^{-1}L_s^{-1}$  is called double inverse Laplace transform with respect to  $p$  and  $s$ . Here we provide double inverse Laplace transform with respect to  $p$  and  $s$  exist for each term in the right hand side of Eqs. (14) and (15).

In order to confirm our method for solving the singular one dimensional pseudo-parabolic fractional equation, we consider the following examples.

**Example 3.1.** Consider the following nonhomogeneous form of a singular one dimensional pseudo-parabolic fractional equation

$$\mathbb{D}_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} - 4, \quad t > 0, \quad x \geq 0 \tag{16}$$

subject to the initial condition

$$u(x, 0) = g(x) = x^2. \tag{17}$$

By applying the above steps, we obtain

$$u_0(x, t) = x^2 + t - 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} \tag{18}$$

and

$$\begin{aligned}
u_{n+1} = & -L_p^{-1}L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial}{\partial x} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right] \\
& - L_p^{-1}L_s^{-1} \left[ \frac{1}{s^\alpha} \int_0^p L_x L_t \left[ \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial}{\partial x} \sum_{n=0}^{\infty} u_n(x, t) \right) \right] dp \right]. \tag{19}
\end{aligned}$$

By Eq.(18) and Eq.(19), we have

$$\begin{aligned} u_1(x, t) &= -L_P^{-1}L_S^{-1} \left[ \frac{1}{S^\alpha} \int_0^P L_x L_t \left[ \frac{\partial}{\partial x}(xu_{0x}) + \frac{\partial^2}{\partial x \partial t}(xu_{0x}) \right] dP \right] \\ &= 4 \frac{t^\alpha}{\Gamma(\alpha + 1)}. \end{aligned}$$

In the same manner, we obtain that

$$u_2 = 0, \quad u_3 = 0, \dots$$

By adding the all terms together, we have

$$u(x, t) = u_0 + u_1 + \dots$$

Therefore, the exact solution is given by

$$u(x, t) = x^2 + t$$

**Example 3.2.** Consider the following nonhomogeneous form of a singular one dimensional pseudo-parabolic fractional equation

$$D_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = x^2 \frac{t^{1-\alpha}}{\Gamma(2-\alpha)}, \quad x \geq 0, \quad t > 0 \quad (20)$$

subject to the initial condition

$$u(x, 0) = g(x) = 0. \quad (21)$$

By applying the above steps, we obtain

$$u_0(x, t) = x^2 t - 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} - 4 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)} \quad (22)$$

$$\begin{aligned} u_1(x, t) &= -L_P^{-1}L_S^{-1} \left[ \frac{1}{S^\alpha} \int_0^P L_x L_t \left[ \frac{\partial}{\partial x}(xu_{0x}) + \frac{\partial^2}{\partial x \partial t}(xu_{0x}) \right] dP \right] \\ &= 4 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)} + 4 \frac{t^\alpha}{\Gamma(\alpha + 1)}. \end{aligned}$$

In the same manner, we obtain that

$$u_2 = 0, u_3 = 0, \dots$$

By adding the all terms together, we have

$$u(x, t) = u_0 + u_1 + \dots$$

Therefore, the exact solution is given by

$$u(x, t) = x^2 t.$$

**Example 3.3.** Consider the following nonhomogeneous form of a singular one dimensional pseudo-parabolic fractional equation

$$D_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = (x^2 + 1) \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} - 8 - 4t, \quad x \geq 0, \quad t > 0 \quad (23)$$

subject to the initial

$$u(x, 0) = g(x) = x^2. \quad (24)$$

By applying the above steps, we obtain

$$\begin{aligned} u_0(x, t) &= x^2 + t - 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} + x^2 t - 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} - 4 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)}. \\ u_1 &= -L_P^{-1} L_S^{-1} \left[ \frac{1}{S^\alpha} \int_0^P L_x L_t \left[ \frac{\partial}{\partial x} (x u_{0x}) + \frac{\partial^2}{\partial x \partial t} (x u_{0x}) \right] dP \right] \\ &= 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} + 4 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)} + 4 \frac{t^\alpha}{\Gamma(\alpha + 1)} \end{aligned}$$

In the same manner, we obtain that

$$u_2 = 0, u_3 = 0, \dots$$

By adding the all terms together, we have

$$u(x, t) = u_0 + u_1 + \dots$$

Therefore, the exact solution is given by

$$u(x, t) = x^2 + t + x^2 t$$



**Example 3.4.** Consider the following nonhomogeneous form of a singular one dimensional pseudo-parabolic fractional equation

$$D_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = x^2 t, \quad x \geq 0, \quad t > 0, \quad (25)$$

subject to the initial condition

$$u(x, 0) = g(x) = 0. \quad (26)$$

By applying the above steps, we obtain

$$\begin{aligned} u_0(x, t) &= x^2 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)}, \\ u_1 &= -L_P^{-1} L_S^{-1} \left[ \frac{1}{S^\alpha} \int_0^P L_x L_t \left[ \frac{\partial}{\partial x} (x u_{0x}) + \frac{\partial^2}{\partial x \partial t} (x u_{0x}) \right] dP \right] \\ &= 4 \frac{t^{2\alpha+1}}{\Gamma(2\alpha + 2)} + 4 \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}. \end{aligned}$$

In the same manner, we obtain that

$$u_2 = 0, \quad u_3 = 0, \dots$$

By adding the all terms together, we have

$$u(x, t) = u_0 + u_1 + \dots$$

Therefore, the exact solution is given by

$$u(x, t) = x^2 \frac{t^{\alpha+1}}{\Gamma(\alpha + 2)} + 4 \frac{t^{2\alpha+1}}{\Gamma(2\alpha + 2)} + 4 \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)}.$$

### 3.2 Convergence Analysis of the method

Finally, we discuss the convergence analysis of the modified double Laplace decomposition methods for the singular nonlinear one dimensional pseudo-parabolic fractional equation which is given by

$$D_t^\alpha u - \frac{1}{x} \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \frac{1}{x} \frac{\partial^2}{\partial x \partial t} \left( x \frac{\partial u}{\partial x} \right) = f(u) \quad (27)$$

Consider the Hilbert space  $H = L_\mu^2((a, b) \times [0, T])$ , defined by the set of applications

$$\left\{ (u, v) : (a, b) \times [0, T] \text{ with } L_s^{-1} L_t^{-1} \left[ \frac{1}{S^\alpha} \int_0^P L_x L_t [u(x, t)](p, s) dp \right] (x, t) < \infty \right\},$$

the scalar product

$$(u, v)_{L^2_\mu(Q)} = \int_Q xu(x, t)v(x, t)dxdt,$$

with the norm

$$\|u\|_H^2 = \int_Q xu^2(x, t)dxdt,$$

where  $Q = (a, b) \times [0, t]$ . Multiplying both sides of Eq. (27) by  $x$ , and we write the equation in the operator form

$$L(u) = xD_t^\alpha u = \frac{\partial u}{\partial x} + x \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial x \partial t} + \frac{\partial^3 u}{\partial x^2 \partial t} - xv \frac{\partial u}{\partial x} + xf(u) \quad (28)$$

For  $L$  is hemicontinuous operator, consider the following hypotheses.

**(H1)**  $(L(u) - L(w), u - w) \geq k\|u - w\|^2$ ,  $k > 0$ ,  $\forall u, w \in H$

**(H2)** Whatever may be  $M > 0$ , there exist a constant  $C(M) > 0$  such that for  $u, w \in H$  with  $\|u\| \leq M$ , and  $\|w\| \leq M$  we have

$$(L(u) - L(w), z_1) \leq C(M)\|u - z_1\| \|w\|$$

for every  $z_1 \in H$ . In the next theorem we follow [1, 8, 10, 11]

**Theorem 3.5.** *(Sufficient condition of convergence) The modified double Laplace decomposition method applied to the singular nonlinear one dimensional pseudo-parabolic fractional equation Eq. (28) without initial and boundary conditions, converges towards a particular solution.*

*Proof.* First, we verify the convergence hypothesis (H1) for the operator  $L(u)$  of Eq. (28). We use the definition of our operator  $L$ , and then we have

$$\begin{aligned} L(u) - L(w) &= \left(\frac{\partial u}{\partial x} - \frac{\partial w}{\partial x}\right) + \left(x \frac{\partial^2 u}{\partial x^2} - x \frac{\partial^2 w}{\partial x^2}\right) + \left(\frac{\partial^2 u}{\partial x \partial t} - \frac{\partial^2 w}{\partial x \partial t}\right) + \left(x \frac{\partial^3 u}{\partial x^2 \partial t}\right) \\ &\quad - \left(xv \frac{\partial u}{\partial x} - xv \frac{\partial w}{\partial x}\right) + x(f(u) - f(w)) \\ &= \frac{\partial}{\partial x}(u - w) + x \frac{\partial^2}{\partial x^2}(u - w) + \frac{\partial^2}{\partial x \partial t}(u - w) \\ &\quad + x \frac{\partial^3}{\partial x^2 \partial t}(u - w) + x(f(u) - f(w)), \end{aligned} \quad (29)$$

Therefore,

$$\begin{aligned}
(L(u) - L(w), u - w) &= \left( \frac{\partial}{\partial x}(u - w), u - w \right) + \left( x \frac{\partial^2}{\partial x^2}(u - w), u - w \right) \\
&+ \left( \frac{\partial^2}{\partial x \partial t}(u - w), u - w \right) + \left( x \frac{\partial^3}{\partial x^2 \partial t}(u - w), u - w \right) \\
&- \left( xv \frac{\partial}{\partial x}(u - w), u - w \right) + (x(f(u) - f(w)), u - w)
\end{aligned} \tag{30}$$

According to the properties of the differential operators  $\frac{\partial}{\partial x}, \frac{\partial^2}{\partial x \partial t}$  in  $H$ , then there exist constants  $\alpha, \beta, \delta, \gamma, k > 0$  such that

$$\left( \frac{\partial}{\partial x}(u - w), u - w \right) \geq \alpha \|u - w\|^2 \tag{31}$$

$$\begin{aligned}
- \left( x \frac{\partial^2}{\partial x^2}(u - w), u - w \right) &\leq |x| \left\| \frac{\partial^2}{\partial x^2}(u - w) \right\| \|u - w\| \\
&\leq b\beta \|u - w\|^2 \\
&\iff
\end{aligned}$$

$$\left( x \frac{\partial^2}{\partial x^2}(u - w), u - w \right) \geq -b\beta \|u - w\|^2, \tag{32}$$

$$\left( x \frac{\partial^2}{\partial x \partial t}(u - w), u - w \right) \geq \gamma \|u - w\|^2, \tag{33}$$

and

$$\begin{aligned}
- \left( x \frac{\partial^3}{\partial x^2 \partial t}(u - w), u - w \right) &\leq |x| \left\| \frac{\partial^3}{\partial x^2 \partial t}(u - w) \right\| \|u - w\| \\
&\leq bk \|u - w\|^2 \\
&\iff
\end{aligned}$$

$$\left( x \frac{\partial^3}{\partial x^2 \partial t}(u - w), u - w \right) \geq -bk \|u - w\|^2. \tag{34}$$

According to the Schwartz inequality, we get

$$\begin{aligned}
\left( xv \frac{\partial}{\partial x}(u - w), u - w \right) &\leq |x| \|v\| \left\| \frac{\partial}{\partial x}(u - w) \right\| \|u - w\| \\
&\leq bM\alpha \|u - w\|^2.
\end{aligned}$$

Hence,

$$- \left( xv \frac{\partial}{\partial x} (u - w), u - w \right) \geq -bM\alpha \|u - w\|^2 \quad (35)$$

By Cauchy Schwartz inequality, where  $\sigma > 0$  and  $f$  is Lipschitzian function, we have

$$\begin{aligned} (-xf(u) - f(w), u - w) &\leq |x| \|f(u) - f(w)\| \|u - w\| \\ &\leq b \|f(u) - f(w)\| \|u - w\| \\ &\leq b\sigma \|u - w\|^2 \\ &\iff \\ (xf(u) - f(w), u - w) &\geq -b\sigma \|u - w\|^2. \end{aligned} \quad (36)$$

Substituting Eq.(31), Eq.(32), Eq.(33), Eq.(34), Eq.(35) and Eq.(36) into equation Eq.(30), gives

$$(L(u) - L(w), u - w) \geq (\alpha - b\beta + \gamma - bk - b\sigma - bM\alpha) \|u - w\|^2$$

$$(L(u) - L(w), u - w) \geq k_1 \|u - w\|^2,$$

where  $k_1 = \alpha - b\beta + \gamma - bk - b\sigma - bM\alpha > 0$ . Hence hypothesis (H1) holds. Now, we verify the convergence hypothesis (H2) for the operators  $L(u)$ . For every  $M > 0$ , there exists a constant  $C(M) > 0$  such that for all  $u, w \in H$  with  $\|u\| \leq M$ , we have

$$(L(u) - L(w), z_1) \leq C(M) \|u - w\| \|z_1\|$$

for every  $z_1 \in H$ . For that we have,

$$\begin{aligned} (L(u) - L(w), z_1) &= \left( \frac{\partial}{\partial x} (u - w), z_1 \right) + \left( x \frac{\partial^2}{\partial x^2} (u - w), z_1 \right) \\ &\quad + \left( \frac{\partial^2}{\partial x \partial t} (u - w), z_1 \right) + \left( x \frac{\partial^3}{\partial x^2 \partial t} (u - w), z_1 \right) \\ &\quad - \left( xv \frac{\partial}{\partial x} (u - w), z_1 \right) + (x(f(u) - f(w)), z_1) \end{aligned}$$

By the Cauchy Schwartz inequality and the fact that  $u$  and  $w$  are bounded, we obtain

$$\left( \frac{\partial}{\partial x} (u - w), z_1 \right) \leq \alpha_1 \|u - w\| \|z_1\|,$$

$$\begin{aligned}
\left(x \frac{\partial^2}{\partial x^2}(u-w), z_1\right) &\leq b\beta_1 \|u-w\| \|z_1\|, \\
\left(\frac{\partial^2}{\partial x \partial t}(u-w), z_1\right) &\leq \gamma_1 \|u-w\| \|z_1\|, \\
\left(x \frac{\partial^3}{\partial x^2 \partial t}(u-w), z_1\right) &\leq bk_1 \|u-w\| \|z_1\|, \\
-\left(xv \frac{\partial}{\partial x}(u-w), z_1\right) &\leq \alpha_1 |x| \|v\| \|u-w\| \|z_1\| \\
&\leq b\alpha_1 M \|u-w\| \|z_1\|, \\
(x(f(u)-f(w)), z_1) &\leq b\sigma_1 \|u-w\| \|z_1\|,
\end{aligned}$$

where the constants  $\alpha_1, \beta_1, \sigma_1, \gamma_1$  and  $k_1$  are positive. Then we have

$$\begin{aligned}
(L(u)-L(w), z_1) &\leq (\alpha_1 + b\beta_1 + \gamma_1 + bk_1 + b\alpha_1 M + b\sigma_1) \|u-w\| \|z_1\| \\
&= C_1(M) \|u-w\| \|z_1\|,
\end{aligned}$$

where  $C_1(M) = \alpha_1 + b\beta_1 + \gamma_1 + bk_1 + b\alpha_1 M + b\sigma_1$ .

□

## 4 Conclusion

In this paper, we have proposed new modified double Laplace decomposition method to solve singular one dimensional pseudo-parabolic fractinal equation. The efficiency and accuracy of the present scheme are validated through examples. This method is much simpler and more efficient in the study of linear and nonlinear PDEs.

## References

- [1] Hassan Eltayeb GADAIN, solving coupled pseudo-parabolic equation using a modified double Laplace decomposition method, *Acta Mathematica Scientia 2018*, **number of volume**(number of issue), vol.38.Ser.B, p. 333-346.
- [2] Francois Dubois, Ana-Cristina Galucio and Nelly point, Introduction of fractional derivative: Theorie and Applications, France, 16 March 2009.

- [3] Abbaoui K and Cherruault Y, Convergence of a domain's method applied to differential equations, *Comput Math Appl*, 1994, 28(5):103-109.
- [4] Abbaoui K and Cherruault Y, Convergence of a domain's method applied to nonlinear equations, *Math Comput Model*, 1994, 20(9):69-73.
- [5] Aminikhah H, An analytical approximation for coupled viscous Burgers equation, *Appl Math Model*, 2013, 37: 59795983.
- [6] Atangana A and Oukouomi Noutchie S C, On multi-Laplace transform for solving nonlinear partial differential equations with mixed derivatives, *Math Probl Eng, Volume 2014*, Article ID 267843, 9 pages.
- [7] Biazar J and Ghazvini H, Exact solution for nonlinear Burgers equation by homotopy perturbation method, *Numer Meth Part D E*, 2009, 25: 833842.
- [8] Do gan Kaya and Inan Ibrahim E, A convergence analysis of the ADM and an application, *Appl Math Comput*, 2005, 161: 10151025.
- [9] Eltayeb H and Kiliman A, A note on solutions of wave, Laplaces and heat equations with convolution terms by using double Laplace transform, *Appl Math Lett*, 2008, 21(12): 13241329.
- [10] Hashim I, Noorani M S M and Said Al-Hadidi M R, Solving the generalized Burgers-Huxley equation using the Adomian decomposition method, *Math Comput Model*, 2006, 43: 14041411.
- [11] Kaya D and Inan I E, A numerical application of the decomposition method for the combined KdV-MKdV equation, *Appl Math Computat*, 2005, 168: 915926.
- [12] Mesloub S, A nonlinear nonlocal mixed problem for a second order pseudoparabolic equation, *J Math Anal Appl*, 2006, 316: 189209.
- [13] Dehghan M, Hamidi A and Shakourifar M, The solution of coupled Burgers equations using Adomian Pad technique, *Appl Math Comput*, 2007, 189: 10341047.
- [14] Kaya D, An explicit solution of coupled viscous Burgers equation by the decomposition method, *IJMMS*, 2001, 27: 675680.

- [15] Kiliman A and Gadain H E, On the applications of Laplace and Sumudu transforms, *J Frankl Inst*, 2010, 347(5): 848862.