

Differential equations and the algebra of confluent spherical functions
on semisimple Lie groups.

Olufemi Opeyemi **OYADARE**

Abstract

We consider the notion of a confluent spherical function on a connected noncompact semisimple Lie group, G , with finite center and of real rank 1, and discuss the properties and relationship of its algebra with the well-known Schwartz algebra of spherical functions on G .

2010 Mathematics Subject Classification: 43A85, 22E30, 22E46

Keywords: Spherical functions: Confluent spherical functions: Poincaré-Birkhoff-Witt Theorem.

Department of Mathematics, Obafemi Awolowo University, Ile-Ife, 220005, Nigeria.

E-mail: *femi_oya@yahoo.com*

§1. Introduction

Let G be a noncompact connected semisimple Lie group with finite center, a maximal compact subgroup K and a Lie algebra \mathfrak{g} having a *Cartan decomposition*,

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}.$$

If we choose a maximal abelian subspace, \mathfrak{a} , of \mathfrak{p} and define $A^+ = \{\exp tH : H \in \mathfrak{a}, t > 0\}$ then G has a *polar decomposition* given as $G = K \cdot cl(A^+) \cdot K$, where $cl(A^+)$ is the closure of A^+ . This decomposition implies that every K -biinvariant function on G is completely determined by its restriction to A^+ . An example of such a function is the (*zonal*) *spherical function*, $\varphi_\lambda, \lambda \in \mathfrak{a}_\mathbb{C}^*$, on G . If we denote the restriction of φ_λ to A^+ as $\tilde{\varphi}_\lambda$, then the following system of differential equations hold:

$$\tilde{q}\tilde{\varphi}_\lambda = \gamma(q)(\lambda)\tilde{\varphi}_\lambda,$$

where $q \in \mathfrak{Q}(\mathfrak{g}_\mathbb{C})(:= U(\mathfrak{g}_\mathbb{C})^K = \text{centralizer of } K \text{ in } U(\mathfrak{g}_\mathbb{C}))$, $\gamma := \gamma_{\mathfrak{g}/\mathfrak{a}}$ is the *Harish-Chandra homomorphism* of $\mathfrak{Q}(\mathfrak{g}_\mathbb{C})$ onto $U(\mathfrak{g}_\mathbb{C})^{\mathfrak{w}}$, the \mathfrak{w} -invariant subspace of $U(\mathfrak{g}_\mathbb{C})$, with \mathfrak{w} denoting the *Weyl group* of the pair $(\mathfrak{g}, \mathfrak{a})$, $tU(\mathfrak{g}_\mathbb{C}) \cap \mathfrak{Q}(\mathfrak{g}_\mathbb{C})$ is the kernel of γ and \tilde{q} is the restriction of q to A^+ . Since

$$\widetilde{q \cdot f} = \tilde{q} \cdot \tilde{f},$$

for every $f \in C^\infty(G//K)$ we conclude that \tilde{q} is the *radial component* of q . We define $q \in \mathfrak{Q}(\mathfrak{g}_\mathbb{C})$ to be *spherical* whenever $q = \tilde{q}$.

The above system of differential equations have been extensively used by Harish-Chandra in the investigation of the nature of the spherical functions, φ_λ , their asymptotic expansions and their contributions to the Schwartz algebras on G . The history of this investigation dated back to the 1950's with the two-volume work of Harish-Chandra, [3(a.)] and [3(b.)], which still attracts the strength of twenty-first century mathematicians (see [10.] and [1.]). Other functions on G satisfying different interesting transformations under members of $\mathfrak{Q}(\mathfrak{g}_\mathbb{C})$ have also been studied in the light of the approach taken by Harish-Chandra. We refer to [5.] and the references cited in it for further discussion.

Now if G is a semisimple Lie group with real rank 1 then it is known (see [3a.]) that the above system of differential equations can be replaced with

$$\delta'(\omega) \cdot \varphi_\lambda = \gamma(\omega)(\lambda) \cdot \varphi_\lambda,$$

where ω is the *Casimir operator* of G and $\delta'(\omega)$ denotes the radial component of the differential operator, $\delta'(\omega)$, associated with ω . If we load the structure of G , as a real rank 1 semisimple Lie group, into the last equation it becomes

$$\left(\frac{d^2}{dt^2} + \{(p+q) \coth t + q \tanh t\} \frac{d}{dt}\right) f_\lambda = \left(\lambda^2 - \frac{(p+2q)^2}{4}\right) f_\lambda,$$

where $p = n(\alpha)$, $q = n(2\alpha)$, $f_\lambda(t) := \varphi_\lambda(\exp tH_0)$ and H_0 is chosen in \mathfrak{a} such that $\alpha(H_0) = 1$ (see [13.], p. 190 for the case of $G = SL(2, \mathbb{R})$). Setting $z = -(\sinh t)^2$ transforms the above ordinary differential equation to the *hypergeometric equation*

$$(z(z-1) \frac{d^2}{dz^2} + ((a+b+1)z - c) \frac{d}{dz} + ab) g_\lambda = 0,$$

where $g_\lambda(z) = f_\lambda(t)$, $z < 0$, $a = \frac{p+2q+2\lambda}{4}$, $b = \frac{p+2q-2\lambda}{4}$ and $c = \frac{p+q+1}{2}$, whose solution is from here given by the *Gauss hypergeometric function*, $F(a, b, c : z)$, defined as

$$F(a, b, c : z) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{z^k}{k!},$$

$|z| < 1$ ([16.], p. 283). It then follows that

$$\varphi_\lambda(\exp tH_0) = F(a, b, c : z),$$

and we conclude that the spherical functions on real rank 1 semisimple Lie groups are essentially the hypergeometric function. In other words, the hypergeometric functions form the spherical functions on any real rank 1 semisimple Lie group.

The confluent hypergeometric function is defined as

$${}_1F(a, c : z) = \lim_{b \rightarrow \infty} F(a, b, c : z/b) = \sum_{k=0}^{\infty} \frac{(a)_k}{(c)_k} \frac{z^k}{k!}.$$

Thus replacing z with z/b in the hypergeometric function, $F(a, b, c : z)$, and computing the limit as $b \rightarrow \infty$ leads to its confluent, ${}_1F(a, c : z)$. Taking the same steps for the above hypergeometric equation shows that ${}_1F(a, c : z)$ satisfies the *confluent differential equation*

$$z \frac{d^2 w}{dz^2} + (c - z) \frac{dw}{dz} - aw = 0.$$

Now since on any real rank 1 semisimple Lie group G , every spherical function is expressible, as seen above, in terms of a hypergeometric function, we refer to ${}_1F(a, c : z)$ as a *confluent spherical function* on G and we denote it by $\varphi_\lambda^\sigma(\exp tH_0)$. i.e.,

$$\varphi_\lambda^\sigma(\exp tH_0) = {}_1F(a, c : z),$$

with $z = -(\sinh t)^2$. It is however noted that if we replace z with z/b in $z = -(\sinh t)^2$, as in the derivation of the confluent hypergeometric equation, then we have

$$t = \sinh^{-1}(i\sqrt{z/b}).$$

So that as $b \rightarrow \infty$ it implies that values of t becomes very small. It then means that the relationship

$$\varphi_\lambda^\sigma(\exp tH_0) = {}_1F(a, c : z),$$

is valid only for sufficiently small values of t . We conclude therefore that the spherical function, $\varphi_\lambda(\exp tH_0)$, becomes a confluent spherical function on G for small values of t .

Our aim in this paper is, therefore, to study the function $\varphi_\lambda(\exp tH_0)$ for small values of t since this corresponds with the study of the confluent spherical function, $\varphi_\lambda^\sigma(\exp tH_0)$ as explained above. In this respect we find the *Stanton-Tomas expansion* of $\varphi_\lambda(\exp tH_0)$ very appropriate to define the general notion of a confluent spherical function.

The paper is arranged as follows: §2. contains a discussion of the radial component of spherical differential operators on any G of arbitrary rank, as discovered by Harish-Chandra ([3a.] and [3b.]), while the motivation for the notion of a confluent spherical function on a real rank 1 semisimple Lie group is developed in §3. This motivation informs our choice of the Stanton-Tomas expansion in the definition of a confluent spherical function. The algebra of these functions are then studied and related with the Schwartz algebra of spherical functions.

An insight into the study of specific confluent spherical functions on the real rank 2 case of $Sp(2, \mathbb{R})$, leading to the consideration of different kinds of Whittaker functions, is contained in Hirano, *et al* [5.]. However the approach taken in this paper is more general than theirs and holds for any real rank 1 semisimple Lie groups, and may be

extended to higher ranks.

§2. Radial Components of Spherical Differential Operators

Consider a noncompact connected real semisimple Lie group G with finite center and with the Lie algebra \mathfrak{g} , whose complexification is denoted as $\mathfrak{g}_{\mathbb{C}}$. We can identify the members of \mathfrak{g} with *left-invariant vector fields* of G in the following manner. For every $X \in \mathfrak{g}$, we define a map

$$X \longmapsto \partial(X),$$

where $\partial(X)$ is to act on members of $C^\infty(G)$ by the requirement

$$(\partial(X)f)(x) := \frac{d}{dt}f(x \cdot \exp tX)|_{t=0}.$$

This depicts $\partial(X)$ as a first order left-invariant differential operator on G associated to every $X \in \mathfrak{g}$ and which satisfies the relation

$$\partial([X, Y]) = \partial(X)\partial(Y) - \partial(Y)\partial(X),$$

for $X, Y \in \mathfrak{g}$. This outlook may be used to introduce left-invariant differential operators of any order on G , by choosing more than one member of \mathfrak{g} at a time. Indeed, if $X_1, \dots, X_r \in \mathfrak{g}$ and we define the map

$$X_1 \cdots X_r \longmapsto \partial(X_1 \cdots X_r)$$

as

$$\partial(X_1 \cdots X_r) = \partial(X_1) \cdots \partial(X_r)$$

then

$$\partial(X_1 \cdots X_r)(x) = \left(\frac{\partial^r}{\partial t_1 \cdots \partial t_r} f(x \cdot \exp tX_1 \cdots \exp tX_r) \right) \Big|_{(t_1, \dots, t_r) = (0, \dots, 0)},$$

which is a left-invariant differential operator on G of order $\leq r$. These operators are analytic and are precisely the endomorphisms of $C^\infty(G)$ generated by $\partial(X)$, $X \in \mathfrak{g}$ ([13], p. 101). Thus if we define $D(G) = \text{span}_{\mathbb{C}}\{\partial(X) : X \in \mathfrak{g}\}$, then $D(G)$ is a subalgebra of the algebra $\text{End}_{\mathbb{C}}(C^\infty(G))$, of all endomorphisms on $C^\infty(G)$, with the identity operator as its identity element.

However, it is known that if $X_1, \dots, X_r \in \mathfrak{g}$, the product $X_1 \cdots X_r$ may not generally be a member of \mathfrak{g} , as may easily be verified with low-dimensional Lie algebras. Thus we should seek a *gadget* in which every product, $X_1 \cdots X_r$, of members of \mathfrak{g} is always found, and then study the structure of the map

$$\partial : X_1 \cdots X_r \longmapsto \partial(X_1 \cdots X_r)$$

with this gadget on the foreground. With this aim in mind we consider the *tensor algebra*, $T(\mathfrak{g}_{\mathbb{C}})$, of the complexification, $\mathfrak{g}_{\mathbb{C}}$, of \mathfrak{g} , given as

$$T(\mathfrak{g}_{\mathbb{C}}) = \mathbb{C} \oplus \mathfrak{g}_{\mathbb{C}} \oplus (\mathfrak{g}_{\mathbb{C}} \otimes \mathfrak{g}_{\mathbb{C}}) \oplus (\mathfrak{g}_{\mathbb{C}} \otimes \mathfrak{g}_{\mathbb{C}} \otimes \mathfrak{g}_{\mathbb{C}}) \oplus \cdots = \bigoplus_{k=0}^{\infty} T^k(\mathfrak{g}_{\mathbb{C}}),$$

where $T^0(\mathfrak{g}_{\mathbb{C}}) := \mathbb{C}$, and $T^k(\mathfrak{g}_{\mathbb{C}}) := \mathfrak{g}_{\mathbb{C}} \otimes \cdots \otimes \mathfrak{g}_{\mathbb{C}}$. $T(\mathfrak{g}_{\mathbb{C}})$ is an associative algebra over \mathbb{C} with identity, and there is a natural map, ι , of $\mathfrak{g}_{\mathbb{C}}$ into $T(\mathfrak{g}_{\mathbb{C}})$ given by identifying $\mathfrak{g}_{\mathbb{C}}$ with the first-order terms ([1.]). $T(\mathfrak{g}_{\mathbb{C}})$ has the following universal property.

2.1 Theorem ([6.], p. 644). If A is any other associative algebra over \mathbb{C} with identity and τ is a linear map of \mathfrak{g} into A , then there exists a unique associative algebra homomorphism $\bar{\tau}$, with $\bar{\tau}(1) = 1$ such that $\bar{\tau} \circ \iota = \tau$. \square

We conclude, from the definitions of ∂ and $T(\mathfrak{g}_{\mathbb{C}})$ above, that

$$\partial : T(\mathfrak{g}_{\mathbb{C}}) \longmapsto D(G).$$

However, as $[X, Y], XY, YX \in T(\mathfrak{g}_{\mathbb{C}})$, for every $X, Y \in \mathfrak{g}_{\mathbb{C}}$ and

$$\partial([X, Y]) - \partial(X)\partial(Y) - \partial(Y)\partial(X) = 0,$$

for every $X, Y \in \mathfrak{g}_{\mathbb{C}}$, it would be necessary to factor, out of $T(\mathfrak{g}_{\mathbb{C}})$, the set generated by all elements of the form $X \otimes Y - Y \otimes X - [X, Y]$, for $X, Y \in \mathfrak{g}_{\mathbb{C}}$. Indeed

$$I = \text{span}_{\mathbb{C}}\{X \otimes Y - Y \otimes X - [X, Y] : X, Y \in \mathfrak{g}_{\mathbb{C}}\}$$

is a two-sided ideal of $T(\mathfrak{g}_{\mathbb{C}})$, and we define

$$U(\mathfrak{g}_{\mathbb{C}}) := T(\mathfrak{g}_{\mathbb{C}})/I.$$

$U(\mathfrak{g}_{\mathbb{C}})$ is also an associative algebra with identity, it contains $\mathfrak{g}_{\mathbb{C}}$ and has the following universal property inherited from $T(\mathfrak{g}_{\mathbb{C}})$.

2.2 Theorem([6.], p. 215). Let ι be the canonical map of $\mathfrak{g}_{\mathbb{C}}$ into $U(\mathfrak{g}_{\mathbb{C}})$, let A be any complex associative algebra with identity, and let φ be a linear mapping of $\mathfrak{g}_{\mathbb{C}}$ into A such that

$$\varphi([X, Y]) = \varphi(X)\varphi(Y) - \varphi(Y)\varphi(X), X, Y \in \mathfrak{g}_{\mathbb{C}}.$$

Then there exists a unique algebra homomorphism $\varphi_0 : U(\mathfrak{g}_{\mathbb{C}}) \longrightarrow A$ with $\varphi_0(1) = 1$ such that $\varphi_0 \circ \iota = \varphi$. \square

The canonical map ι , in the above theorem is one-to-one ([6.], p. 217) and the object $U(\mathfrak{g}_{\mathbb{C}})$ is called the *universal enveloping algebra* of $\mathfrak{g}_{\mathbb{C}}$. One of the most fundamental results in the theory of $U(\mathfrak{g}_{\mathbb{C}})$ is the following, which gives a concrete way of constructing it.

2.3 Theorem(Poincaré-Birkhoff-Witt Theorem)([1.], p. 32). If X_1, \dots, X_n is a basis of $\mathfrak{g}_{\mathbb{C}}$ over \mathbb{C} , then the monomials $\iota(X_1)^{j_1} \cdots \iota(X_n)^{j_n}$, $j_k \geq 0, k = 1, \dots, n$, form a basis of $U(\mathfrak{g}_{\mathbb{C}})$ over \mathbb{C} . \square

The inclusion of ι in the above is not necessary since it is a one-to-one map. However, the members of $U(\mathfrak{g}_{\mathbb{C}})$ may seem to be difficult to handle if we only the definition $U(\mathfrak{g}_{\mathbb{C}}) := T(\mathfrak{g}_{\mathbb{C}})/I$ in mind. However recalling, from Theorem 2.2, that, with $A = D(G)$, $\partial : \mathfrak{g}_{\mathbb{C}} \longrightarrow D(G)$ is a (natural) homomorphism such that $\partial([X, Y]) = \partial(X)\partial(Y) - \partial(Y)\partial(X), X, Y \in \mathfrak{g}_{\mathbb{C}}$, and which also extends to all of $U(\mathfrak{g}_{\mathbb{C}})$, this implies that the members of $U(\mathfrak{g}_{\mathbb{C}})$ are more concrete than predicted by the Poincaré-Birkhoff-Witt Theorem. Indeed we have the following major result that gives a different outlook on $U(\mathfrak{g}_{\mathbb{C}})$.

2.4 Theorem([1.], p. 32). The algebra homomorphism $\partial : U(\mathfrak{g}_{\mathbb{C}}) \longrightarrow D(G)$ is an algebra isomorphism onto. \square

The message of these theorems is that the members of $U(\mathfrak{g}_{\mathbb{C}})$ are *mixed derivatives*, so that $U(\mathfrak{g}_{\mathbb{C}})$ is the algebra of all left-invariant differential operators on G . This allows us to study $U(\mathfrak{g}_{\mathbb{C}})$ as the *house* of all left-invariant differential operators on $C^\infty(G)$. Furthermore $U(\mathfrak{g}_{\mathbb{C}})$ may also be realized as the algebra of *right-invariant* differential operators on G via the anti-isomorphism ∂_r given as

$$(\partial_r(X_1 \cdots X_r)f)(x) = \left(\frac{\partial^r}{\partial t_1 \cdots \partial t_r} f(\exp t_1 X_1 \cdots \exp t_r X_r \cdot x) \right) \Big|_{(t_1, \dots, t_r) = (0, \dots, 0)}.$$

This second realization of $U(\mathfrak{g}_{\mathbb{C}})$ suggests that there are some of its members which

are both left- and right-invariant. i.e., members $q \in U(\mathfrak{g}_{\mathbb{C}})$ in which $qX = Xq$, for all $X \in \mathfrak{g}_{\mathbb{C}}$. This set of members that are both left- and right-invariant is the *center* of $U(\mathfrak{g}_{\mathbb{C}})$ and is denoted by $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$. Though the algebra $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ is abelian and sufficient in the harmonic analysis on G , we shall however consider the larger subalgebra \mathfrak{Q} of $U(\mathfrak{g}_{\mathbb{C}})$ defined as the centralizer of K in $U(\mathfrak{g}_{\mathbb{C}})$. i.e.,

$$\mathfrak{Q} = \{q \in U(\mathfrak{g}_{\mathbb{C}}) : Ad(k)q = q, k \in K\}.$$

This is due to the fact that we are ultimately interested in the study of K -biinvariant functions on G . It is the *radial component* of members of \mathfrak{Q} , viewed as a subalgebra of the algebra, $D(G)$, of left-invariant differential operators on G , that we set out to compute in this section. This is reminiscence of the classical method of finding the *normal form* of an ordinary differential operator. However we need to have a generalization of the *polar decomposition of matrices* to members of G in order to start. We take a cue from the example of the case $G = SL(2, \mathbb{R})$, where the generalization of *polar coordinates* and *normal form* are easily seen.

Let $G = SL(2, \mathbb{R}) = \{x = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL(2, \mathbb{R}) : ad - bc = 1\}$ with Lie algebra $\mathfrak{g} = sl(2, \mathbb{R}) = \{X \in GL(2, \mathbb{R}) : tr(X) = 0\}$ and complexification $\mathfrak{g}_{\mathbb{C}} = sl(2, \mathbb{C}) = \{X \in GL(2, \mathbb{C}) : tr(X) = 0\}$. The matrices $H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, $X = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, and

$Y = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ are members of \mathfrak{g} and are such that the set $\{X - Y, H, X + Y\}$ form

a basis for \mathfrak{g} . Now since, for $t \in \mathbb{R}$, $\exp tH = \begin{pmatrix} e^t & 0 \\ 0 & e^{-t} \end{pmatrix} =: a_t$ and, for $\theta \in [0, 2\pi]$,

$\exp \theta(X - Y) = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} =: u_{\theta}$, form the closed subgroups A and K of G , the

basis of \mathfrak{g} above implies that $G = K \cdot A \cdot K$. More precisely we have $G = K \cdot cl(A^+) \cdot K$, where $cl(A^+)$ stands for the closure of $\{a_t : t > 0\}$. This is the polar decomposition which is known to generalize to any noncompact connected semisimple Lie group, with finite center. Another way to establish the polar decomposition, which easily extends

to more general semisimple Lie groups, is by considering the map

$$\varphi : K \times cl(A^+) \times K \longrightarrow G : (k_{\theta_1}, a_t, k_{\theta_2}) \longmapsto \varphi(k_{\theta_1}, a_t, k_{\theta_2}) = k_{\theta_1} a_t k_{\theta_2}.$$

This map is a diffeomorphism onto, and we can seek its differentials on the basis elements $X - Y, H, X + Y$ of \mathfrak{g} . Indeed, since the tangent spaces to K and A at 1 are $\mathfrak{k} = \{\theta(X - Y) : \theta \in \mathbb{R}\} =: \mathbb{R}(X - Y)$ and $\mathfrak{a} = \{tH : t \in \mathbb{R}\} =: \mathbb{R}H$, then we have that

$$\frac{\partial}{\partial x_1} = \frac{\partial}{\partial \theta_1}, \quad \frac{\partial}{\partial x_2} = \frac{\partial}{\partial t}, \quad \frac{\partial}{\partial x_3} = \frac{\partial}{\partial \theta_2},$$

in the formula

$$\partial\varphi\left(\frac{\partial}{\partial x_j}\Big|_m\right) = \sum_{k=1}^3 \frac{\partial(y_k \circ \varphi)}{\partial x_j}\Big|_m \cdot \frac{\partial}{\partial y_k}\Big|_{\varphi(m)},$$

where (x_1, x_2, x_3) and (y_1, y_2, y_3) are coordinate systems about m and $\varphi(m)$, respectively (cf. [14.], p. 17). Since, in this case, $m = 1$, we have $\varphi(m) = 1$. Thus, as the tangent space to G at $\varphi(1) = 1$ is \mathfrak{g} , with an orthonormal basis $\{X - Y, H, X + Y\}$, we also have that

$$\frac{\partial}{\partial y_1} = (X - Y), \quad \frac{\partial}{\partial y_2} = H, \quad \frac{\partial}{\partial y_3} = (X + Y).$$

Hence

$$\begin{aligned} & \frac{\partial(y_1 \circ \varphi)}{\partial x_1}\Big|_1 \\ &= \frac{\partial(y_1 \circ \varphi)}{\partial \theta_1}\Big|_{(\theta_1, t, \theta_2)=(0,0,0)} \\ &= \frac{\partial}{\partial \theta_1}(\exp \theta_1(X - Y) \cdot a_t \cdot k_{\theta_2})\Big|_{(\theta_1, t, \theta_2)=(0,0,0)} \\ &= \frac{\partial}{\partial \theta_1}(a_t k_{\theta_2} \exp \theta_1(X - Y)^{k(-\theta_2)a(-t)})\Big|_{(\theta_1, t, \theta_2)=(0,0,0)} \\ & \hspace{15em} (\text{since } \exp(Z)y = y \exp(Zy^{-1}) \text{ for } Z \in \mathfrak{g}, y \in G.) \\ &= k_0 a_0 \frac{d}{d\theta_1}(\exp \theta_1(X - Y)^{k(-\theta_2)a(-t)})\Big|_{\theta_1=0} \\ &= (X - Y)^{k(-\theta_2)a(-t)} \\ &= k_{(-\theta_2)} \cdot a_{(-t)} \cdot (X - Y) \cdot a_t \cdot k_{\theta_2} \\ &= (\cosh 2t)(X - Y) + (\sin 2\theta_2 \sinh 2t)H + (-\cos 2\theta_2 \sinh 2t)(X + Y) \end{aligned}$$

In the same manner

$$\begin{aligned}\frac{\partial(y_2 \circ \varphi)}{\partial x_2} \Big|_1 &= H^{k(-\theta_2)} = 0(X - Y) + (\cos 2\theta_2)H + (\sin 2\theta_2)(X + Y), \\ \frac{\partial(y_3 \circ \varphi)}{\partial x_3} \Big|_1 &= (X - Y) = 1(X - Y) + 0H + 0(X + Y).\end{aligned}$$

We then have the specifications of the differential, $d\varphi$, of φ on the basis elements, $\frac{\partial}{\partial\theta_1}$, $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial\theta_2}$, as

$$d\varphi : \begin{cases} \frac{\partial}{\partial\theta_1} \mapsto (\cosh 2t)(X - Y) + (\sin 2\theta_2 \sinh 2t)H + (-\cos 2\theta_2 \sinh 2t)(X + Y) \\ \frac{\partial}{\partial t} \mapsto 0(X - Y) + (\cos 2\theta_2)H + (\sin 2\theta_2)(X + Y) \\ \frac{\partial}{\partial\theta_2} \mapsto 1(X - Y) + 0H + 0(X + Y) \end{cases}$$

The *Jacobian* of this transformation is then $\sinh 2t$, so that the corresponding Haar measure, dG , of G is also $dG = \frac{1}{2} \sinh 2t d\theta_1 dt d\theta_2$. Its inverse, $(d\varphi)^{-1}$, is then given as

$$(d\varphi)^{-1} : \begin{cases} (X - Y) \mapsto \frac{\partial}{\partial\theta_2} \\ H \mapsto \cos 2\theta_2 \frac{\partial}{\partial t} + \frac{\sin 2\theta_2}{\sinh 2t} \left(\frac{\partial}{\partial\theta_1} - \cosh 2t \frac{\partial}{\partial\theta_2} \right) \\ (X + Y) \mapsto \sin 2\theta_2 \frac{\partial}{\partial t} - \frac{\cos 2\theta_2}{\sinh 2t} \left(\frac{\partial}{\partial\theta_1} - \cosh 2t \frac{\partial}{\partial\theta_2} \right) \end{cases}$$

by simple substitution of the terms for $\frac{\partial}{\partial\theta_2}$, $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial\theta_1}$, respectively.

Now, since the above expression for $(d\varphi)^{-1}$ imply that $d\varphi$ is bijective everywhere on $K \times A^+ \times K$ it follows that ([13.], p. 190) that any analytic differential operator D on $G^+ := KA^+K$ gives rise to a *unique* differential operator D^φ on $K \times A^+ \times K$, called the *polar form of D* , such that, for any $f \in C^\infty(G^+)$, we have

$$(Df) \circ \varphi = D^\varphi(f \circ \varphi).$$

The *composition with φ* in this equation means *restriction to G^+* . If we now denote the restriction of f to A^+ by \tilde{f} , then the last equation above becomes

$$\widetilde{Df} = \tilde{D}\tilde{f}$$

This means that \tilde{D} is the *radial component* of the differential operator D on G^+ , whose existence is proved in [3a.], p. 265, and is called *spherical* whenever $D = \tilde{D}$. Now since

$D(G) \cong U(\mathfrak{g}_{\mathbb{C}})$ it is sufficient to consider D^φ for $D = Z \in \mathfrak{g}_{\mathbb{C}} = \mathfrak{sl}(2, \mathbb{C})$. Indeed, for every $D = Z \in \mathfrak{g}_{\mathbb{C}}$, we have that, $D^\varphi = (d\varphi)^{-1}(Z)$ and since a standard basis $\{H', X', Y'\}$ of $\mathfrak{g}_{\mathbb{C}}$ is given as

$$H' = -i(X - Y), X' = \frac{1}{2}(H + i(X + Y)), Y' = \frac{1}{2}(H - i(X + Y)),$$

we have that

$$H'^\varphi = (d\varphi)^{-1}(H') = -i \frac{\partial}{\partial \theta_2},$$

$$X'^\varphi = (d\varphi)^{-1}(X') = \frac{ie^{-2i\theta_2}}{\sinh 2t} \frac{\partial}{\partial \theta_1} + e^{-2i\theta_2} \frac{\partial}{\partial t} - i \frac{\cosh 2t}{\sinh 2t} e^{-2i\theta_2} \frac{\partial}{\partial \theta_2},$$

and

$$Y'^\varphi = (d\varphi)^{-1}(Y') = \frac{-ie^{2i\theta_2}}{\sinh 2t} \frac{\partial}{\partial \theta_1} + e^{2i\theta_2} \frac{\partial}{\partial t} + i \frac{\cosh 2t}{\sinh 2t} e^{2i\theta_2} \frac{\partial}{\partial \theta_2}.$$

The Poincaré-Birkhoff-Witt Theorem above then implies that we consider the radial components of the monomials given as

$$H'^{j_1} X'^{j_2} Y'^{j_3}, j_1, j_2, j_3 \geq 0,$$

in order to exhaust the members of $U(\mathfrak{g}_{\mathbb{C}})$. It is sufficient, for a start, however to consider the center $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ or the *centralizer*, $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, of K in $U(\mathfrak{g}_{\mathbb{C}})$, both of which are commutative subalgebras. Indeed, $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ is the commutative polynomial algebra in the single variable

$$\omega = (H')^2 + 2H' + 4Y'X',$$

the *Casimir operator* of G . i.e., $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}}) = \mathbb{C}[\omega]$ ([8.], p.195), while $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$ is the commutative polynomial algebra in the two variables ω and $X - Y$. i.e., $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}}) = \mathbb{C}[\omega, X - Y]$ ([8.], p.196). Clearly $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}}) \subset \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$. We however use the *normalized casimir operator*, ω' , given as

$$\omega' = (H')^2 + 2H' + 4Y'X' + 1,$$

so that

$$\omega'^\varphi = (d\varphi)^{-1}(\omega') = \left[\frac{1}{\sinh^2(2t)} \left(\frac{\partial^2}{\partial \theta_1^2} + \frac{\partial^2}{\partial \theta_2^2} \right) - 2 \frac{\cosh 2t}{\sinh^2(2t)} \frac{\partial^2}{\partial \theta_1 \partial \theta_2} \right] + \left[\frac{\partial^2}{\partial t^2} + 2 \frac{\cosh 2t}{\sinh 2t} + 1 \right]$$

is the restriction of ω' to G^+ , and may be referred to as the polar form of ω' . This shows that members of $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$, indeed of $U(\mathfrak{g}_{\mathbb{C}})$, are essentially partial differential operators on

G^+ . Hence the radial component, $\delta'(\omega')$, of the normalized Casimir operator, ω' , which is the restriction of the above ω' to A^+ , is simply

$$\delta'(\omega') = \frac{d^2}{dt^2} + 2 \frac{\cosh 2t}{\sinh 2t} \frac{d}{dt} + 1,$$

(see also [7.], p. 73) reducing the mixed derivatives from $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ to the ordinary derivatives $\frac{d^r}{dt^r}$, $0 \leq r \leq 2$, on spherical functions on G .

The following well-known result of [8.], p. 199, explains that the eigenfunctions of $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$, or of ω' , are exactly the spherical functions on G .

2.5 Theorem. A K -biinvariant C^∞ function f on G , with $f(1) = 1$, is a spherical function iff $\omega' = \lambda^2 f$ for some $\lambda \in \mathbb{C}$. \square

A detailed proof of Theorem 2.5 is contained in [9.], p. 88. Now as every element $q \in U(\mathfrak{sl}(2, \mathbb{C}))$, of degree $\leq r$, may be written as

$$q = \gamma H^r + \sum_{l+m+n \leq r, m \leq r-1} \beta_{l,m,n} (X - Y)^l H^m (X + Y)^n,$$

with where γ and $\beta_{l,m,n}$ are constants, we may generalize the above expression for $\delta'(\omega')$ to all members of $U(\mathfrak{sl}(2, \mathbb{C}))$ in the following manner. Let \mathcal{R} be the complex algebra of functions on $(0, \infty)$ that are generated by $(\sinh 2t)^{-1}$ and $\cosh 2t \cdot (\sinh 2t)^{-1}$. We know that $(\frac{d}{dt})\mathcal{R} \subset \mathcal{R}$, since the derivatives of the generators are all in \mathcal{R} . The reduction of every $q \in U(\mathfrak{sl}(2, \mathbb{C}))$, and not just of $\mathfrak{Z}(\mathfrak{sl}(2, \mathbb{C}))$, to ordinary derivatives is established using the above expression for q and the method of using $(d\varphi)^{-1}$ in the calculation of D^φ , for every $D = Z \in \mathfrak{sl}(2, \mathbb{C})$, as enumerated for ω' above.

2.6 Proposition([13.], p. 238). If $q \in U(\mathfrak{sl}(2, \mathbb{C}))$ is of degree $\leq r$, then there exist $f_0, \dots, f_{r-1} \in \mathbb{C} \cdot 1 \otimes \mathcal{R}$ such that, for any $\varphi \in C^\infty(G//K)$, we have

$$(q \cdot \varphi) = (\delta'(q) \cdot \varphi)$$

on A^+ , where the operator $\delta'(q)$ is given as

$$\delta'(q) = \gamma \frac{d^r}{dt^r} + \sum_{0 \leq j \leq r-1} f_j \cdot \frac{d^j}{dt^j},$$

for some constant γ . \square

If $r = 2$ and $q = \omega' \in \mathfrak{Z}(\mathfrak{sl}(2, \mathbb{C}))$, the conclusion of Proposition 2.6 implies that

$$\delta'(\omega') = \gamma \frac{d^2}{dt^2} + f_1 \frac{d}{dt} + f_0,$$

which is in conformity with the direct computations above, where we see that γ is 1, $f_1(t)$ is $2\frac{\cosh 2t}{\sinh 2t}$ and $f_0(t)$ is 1. The operator $\delta'(q)$ in Proposition 2.6 above may then be called the radial component of every $q \in U(\mathfrak{sl}(2, \mathbb{C}))$. It would be a huge step to generalize this Proposition to every q in the universal enveloping algebra, $U(\mathfrak{g}_{\mathbb{C}})$, of the complexification, $\mathfrak{g}_{\mathbb{C}}$, of a real connected semisimple Lie algebra, \mathfrak{g} , of G . To this end we extract the basic features in the above case of $G = SL(2, \mathbb{R})$ as follows:

(i.) Computation of the differential(s) of the *Cartan decomposition map*, $K \times A^+ \times K \longrightarrow G^+$.

(ii.) Use this differential, in (i.), to find the radial component for every member of $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ or of $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$.

Though the programme to solve items (i.) and (ii.) above may not be as straightforward as we have seen for $\mathfrak{g}_{\mathbb{C}} = \mathfrak{sl}(2, \mathbb{C})$, the following result sets in motion the process of dealing with (i.). To this end, let $\varphi : K \times A \times K \longrightarrow G$ and $G^+ = KA^+K$.

2.7 Proposition([2.], p. 125). The map $\varphi : K \times A \times K \longrightarrow G$, given as $\varphi(k_1, h, k_2) := k_1 h k_2$, is submersive on $K \times A^+ \times K$. In particular, G^+ is open in G and φ is an open map of $K \times A^+ \times K$ onto G^+ .

Proof. We prove that the differential $(d\varphi)_{(k_1, h, k_2)} =: D$ maps $\mathfrak{t} \times \mathfrak{a} \times \mathfrak{t}$ onto \mathfrak{g} . To this end, let $Z_1, Z_2 \in \mathfrak{t}, R \in \mathfrak{a}$, then

$$\begin{aligned}
(d\varphi)_{(k_1, h, k_2)}(Z_1, R, Z_2) &= D(Z_1, R, Z_2) \\
&= D(Z_1, 0, 0) + D(0, R, 0) + D(0, 0, Z_2) \\
&= (d\varphi)_{(k_1, h, k_2)}(Z_1, 0, 0) + (d\varphi)_{(k_1, h, k_2)}(0, R, 0) + (d\varphi)_{(k_1, h, k_2)}(0, 0, Z_2) \\
&= \left(\frac{d}{d\theta_1} \varphi(k_1 \exp \theta_1 Z_1 \cdot h \exp 0 \cdot k_2 \exp 0)\right)_{|\theta_1=0} \\
&\quad + \left(\frac{d}{dt} \varphi(k_1 \exp 0 \cdot h \exp tR \cdot k_2 \exp 0)\right)_{|t=0} \\
&\quad + \left(\frac{d}{d\theta_2} \varphi(k_1 \exp 0 \cdot h \exp 0 \cdot k_2 \exp \theta_2 Z_2)\right)_{|\theta_2=0} \\
&= \left(\frac{d}{d\theta_1} \varphi(k_1 \exp \theta_1 Z_1 \cdot h k_2)\right)_{|\theta_1=0} + \left(\frac{d}{dt} \varphi(k_1 \cdot h \exp tR \cdot k_2)\right)_{|t=0} \\
&\quad + \left(\frac{d}{d\theta_2} \varphi(k_1 h \cdot k_2 \exp \theta_2 Z_2)\right)_{|\theta_2=0}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{d}{d\theta_1} \varphi(k_1 k_2 h \cdot \exp \theta_1 Z_1^{(hk_2)^{-1}}) \right) \Big|_{\theta_1=0} + \left(\frac{d}{dt} \varphi(k_1 h k_2 \cdot \exp t R^{k_2^{-1}}) \right) \Big|_{t=0} \\
&+ \left(\frac{d}{d\theta_2} \varphi(k_1 h k_2 \cdot \exp \theta_2 Z_2) \right) \Big|_{\theta_2=0} \\
&= Z^{(hk_2)^{-1}} + R^{k_2^{-1}} + Z_2 \\
&= Ad(k_2^{-1}) \cdot (Z_1^{h^{-1}}) + Ad(k_2^{-1}) \cdot R + Ad(k_2^{-1}) \cdot (Z_2^{k_2}) \\
&= Ad(k_2^{-1}) \cdot (Z_1^{h^{-1}} + R + Z_2^{k_2})
\end{aligned}$$

Hence $(Ad(k_2) \circ D)(Z_1, R, Z_2) = Z_1^{h^{-1}} + R + Z_2^{k_2}$, showing that D is one-to-one. The surjectivity of D would hold if we show that $\mathfrak{g} = \mathfrak{t}^{h^{-1}} + \mathfrak{a} + \mathfrak{t}$ for every $h \in A^+$. Indeed, it is sufficient to verify that $\theta \mathfrak{n}$ is contained in $\mathfrak{t}^{h^{-1}} + \mathfrak{a} + \mathfrak{t}$, where θ is a *Cartan involution* on \mathfrak{g} , $\mathfrak{n} = \sum_{\alpha \in \Delta^+} \mathfrak{g}_\alpha$, and Δ^+ is a set of *positive restricted roots* of $(\mathfrak{g}, \mathfrak{a})$. To this end, let $X \in \mathfrak{g}_\alpha, \alpha \in \Delta^+$, then

$$(X + \theta X)^{h^{-1}} = e^{-\alpha(\log h)} X + e^{\alpha(\log h)} (\theta X),$$

so that

$$\begin{aligned}
(X + \theta X)^{h^{-1}} - e^{-\alpha(\log h)} (X + \theta X) &= e^{-\alpha(\log h)} X + e^{\alpha(\log h)} (\theta X) - e^{-\alpha(\log h)} X - e^{-\alpha(\log h)} (\theta X) \\
&= (e^{\alpha(\log h)} - e^{-\alpha(\log h)}) (\theta X).
\end{aligned}$$

i.e.,

$$\begin{aligned}
\theta X &= (e^{\alpha(\log h)} - e^{-\alpha(\log h)})^{-1} ((X + \theta X)^{h^{-1}} - e^{-\alpha(\log h)} (X + \theta X)) \\
&= (e^{\alpha(\log h)} - e^{-\alpha(\log h)})^{-1} (X + \theta X)^{h^{-1}} - (e^{\alpha(\log h)} - e^{-\alpha(\log h)})^{-1} e^{-\alpha(\log h)} (X + \theta X) \dots (*) \\
&\in \mathfrak{t}^{h^{-1}} + \mathfrak{t} && \text{(since } (X + \theta X) \in \mathfrak{t} \text{)} \\
&\subset \mathfrak{t}^{h^{-1}} + \mathfrak{a} + \mathfrak{t}.
\end{aligned}$$

This ends the proof as expected. \square

We see that $(d\varphi)_{(1,h,1)} = Z_1^{h^{-1}} + R + Z_2$, in anticipation of its use on the K -biinvariant functions on G . The proof of Proposition 2.7 above gives a formula for the

first differential, $(d\varphi)_{(k_1, h, k_2)}$, of φ . However since we are ultimately interested in the *radial component* of an arbitrary C^∞ spherical differential operator on G , which may have second, third, and higher derivatives, we compute higher order derivatives of φ to give the full differential, which we shall denote by $(d\varphi)_{(k_1, h, k_2)}^\infty$. This is mainly because the property of $\mathfrak{Z}(sl(2, \mathbb{C}))$, as a polynomial algebra in the variable ω' , has not been found generalizable to arbitrary semisimple G and \mathfrak{g} . Indeed we have the following.

2.8 Proposition ([2.], p. 127). The full differential, $(d\varphi)_{(k_1, h, k_2)}^\infty$, of the map φ in Proposition 2.7 is given on $U(\mathfrak{t}_{\mathbb{C}}) \oplus U(\mathfrak{a}_{\mathbb{C}}) \oplus U(\mathfrak{t}_{\mathbb{C}})$ as

$$(d\varphi)_{(k_1, h, k_2)}^\infty(\xi_1 \otimes u \otimes \xi_2) = \xi_1^{(hk_2)^{-1}} u^{k_2^{-1}} \xi_2,$$

where $\xi_1, \xi_2 \in U(\mathfrak{t}_{\mathbb{C}})$ and $u \in U(\mathfrak{a}_{\mathbb{C}})$. In particular

$$(d\varphi)_{(1, h, 1)}^\infty(\xi_1 \otimes u \otimes \xi_2) = \xi_1^{h^{-1}} u \xi_2. \quad \square$$

Just as in Proposition 2.7, the map

$$D_h : U(\mathfrak{t}_{\mathbb{C}}) \oplus U(\mathfrak{a}_{\mathbb{C}}) \oplus U(\mathfrak{t}_{\mathbb{C}}) \longrightarrow U(\mathfrak{g}_{\mathbb{C}})$$

defined as $D_h = (d\varphi)_{(1, h, 1)}^\infty$ is surjective. Thus for every $q \in U(\mathfrak{g}_{\mathbb{C}})$ there exists $\tau_h \in U(\mathfrak{t}_{\mathbb{C}}) \oplus U(\mathfrak{a}_{\mathbb{C}}) \oplus U(\mathfrak{t}_{\mathbb{C}})$ such that $D_h(\tau_h) = q$. If we assume that τ_h depends smoothly on h , then the map $h \mapsto \tau_h$ leads to a differential operator on $K \times A^+ \times K$ which, at the points of $(1) \times A^+ \times (1)$, is simply q expressed in polar coordinates. To find the formula for τ_h we proceed as follows.

Let $q \mapsto t(q)$ be the projection of $U(\mathfrak{g}_{\mathbb{C}})$ onto $U(\mathfrak{a}_{\mathbb{C}})U(\mathfrak{t}_{\mathbb{C}})$, which corresponds to the direct sum $U(\mathfrak{g}_{\mathbb{C}}) = U(\mathfrak{a}_{\mathbb{C}})U(\mathfrak{t}_{\mathbb{C}}) \oplus \theta(\mathfrak{n})U(\mathfrak{g}_{\mathbb{C}})$, then $\deg(t(q)) \leq \deg(q)$. In view of the f'_j s in Proposition 2.6, we define f_α and $g_\alpha, \alpha \in \Delta^+$, on A^+ by $f_\alpha = (\xi_\alpha - \xi_{-\alpha})^{-1}$ and $g_\alpha = \xi_{-\alpha}(\xi_\alpha - \xi_{-\alpha})^{-1}$, respectively, where $\xi_\lambda = e^{\lambda \log}, \lambda \in \mathfrak{a}_{\mathbb{C}}$. Also, let \mathcal{R}_0 be the algebra with unit generated over, \mathbb{C} , by the f_α and g_α , and, for any integer $d \geq 1$, let $\mathcal{R}_{0,d}$ be the linear span of the monomials in these generators of degree d . Now if we put

$$\mathcal{R}_0^+ = \sum_{d \geq 1} \mathcal{R}_{0,d},$$

then for every $q \in U(\mathfrak{g}_{\mathbb{C}})$ of degree m , there exist $\xi_i, \xi'_i \in U(\mathfrak{t}_{\mathbb{C}}), u_i \in U(\mathfrak{a}_{\mathbb{C}})$ and $\varphi_i \in \mathcal{R}_0^+, 1 \leq i \leq n$ such that

$$q = t(q) + \sum_{1 \leq i \leq n} \varphi_i(h) \xi_i^{h^{-1}} u_i \xi'_i, \quad h \in A^+,$$

and we may take

$$\tau_h = 1 \otimes t(q) + \sum_{1 \leq i \leq n} \varphi_i(h) \xi_i \otimes u_i \otimes \xi'_i, h \in A^+, q \in U(\mathfrak{a}_{\mathbb{C}}).$$

This is the appropriate generalization of the expression for q in $U(\mathfrak{sl}(2, \mathbb{C}))$ of degree $\leq r$ as given after Theorem 2.5 above. This direct comparison with the case of $\mathfrak{sl}(2, \mathbb{C})$ implies that we need to use the general expression for q above to seek the generalization of Proposition 2.6 to all members of $U(\mathfrak{g}_{\mathbb{C}})$ for any semisimple Lie algebra \mathfrak{g} . It is sufficient, in our present case, to seek this generalization to all members of $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$ as we now do next. First a little preparation.

Take $\beta_{\mathfrak{n}} : U(\mathfrak{g}_{\mathbb{C}}) \rightarrow U(\mathfrak{a}_{\mathbb{C}})$ be the projection corresponding to the direct sum $U(\mathfrak{g}_{\mathbb{C}}) = U(\mathfrak{a}_{\mathbb{C}}) \oplus (\mathfrak{t}U(\mathfrak{g}_{\mathbb{C}}) + U(\mathfrak{g}_{\mathbb{C}})\mathfrak{n})$, having $\mathfrak{Q} \cap (\mathfrak{t}U(\mathfrak{g}_{\mathbb{C}}))$ as its kernel ([3(a.)], p. 260) and define a map $\gamma_{\mathfrak{n}} : \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}}) \rightarrow U(\mathfrak{a}_{\mathbb{C}})$ by the specification

$$\gamma_{\mathfrak{n}}(q)(\lambda) = \beta_{\mathfrak{n}}(q)(\lambda - \rho), \lambda \in \mathfrak{a}_{\mathbb{C}}^*, q \in \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}}), \rho = \frac{1}{2} \sum_{\alpha \in \Delta^+} \dim(\mathfrak{g}_{\alpha}) \cdot \alpha.$$

$\gamma_{\mathfrak{n}}$ is a homomorphism ([3(a.)], p. 260), is independent of the choice of \mathfrak{n} and is called the *Harish-Chandra homomorphism*. We denote it simply as γ which, for $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R})$, is given on $\omega' = (H')^2 + 2H' + 4Y'X'$ as $\gamma(\omega') = H^2 \mapsto \frac{d^2}{dt^2}$. (see [7.], p. 51 and use the isomorphism in Theorem 2.4) We state the major result of this section.

2.9 Theorem([2.], p. 129 and [3(a.)], p. 267). Given any analytic spherical differential operator E on G^+ there is a unique analytic differential operator \tilde{E} on A^+ , the radial component of E , such that $\tilde{E}f = \tilde{E}f, f \in C^\infty(G//K)$. The map $E \mapsto \tilde{E}$ is a homomorphism that does not increase degree. If $E = q \in \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, then the radial component, written as $\delta'(q)$, is given as

$$\delta'(q) = e^{-\rho} \cdot \gamma(q) \cdot e^{\rho} + \sum_{1 \leq i \leq n} \varphi_i u_i,$$

where $u_i \in U(\mathfrak{g}_{\mathbb{C}})$, $\varphi_i \in \mathcal{R}_0^+$ and $\deg(u_i) < \deg(q)$. \square

It is clear that Theorem 2.9 generalizes the assertions of Proposition 2.6, at least to all members of $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, and it sets the stage for analysis of the differential equations satisfied by spherical functions on G . Indeed, by Theorem 2.5 we have that $\omega' \varphi = \lambda^2 \varphi$ which, when combined with Proposition 2.9, (see also Lemma 23 of [3(a.)]) implies that

$\delta'(\omega')\varphi = \lambda^2\varphi$ where, according to Theorem 2.9,

$$\delta'(\omega') = e^{-\rho} \cdot \gamma(\omega') \cdot e^{\rho} + \sum_{1 \leq i \leq n} \varphi_i u_i,$$

and ω' is the *normalized* Casimir operator of $U(\mathfrak{g}_{\mathbb{C}})$. Even though the relation $\mathfrak{Z}(sl(2, \mathbb{C})) = \mathbb{C}[\omega']$ does not generalize to arbitrary $\mathfrak{g}_{\mathbb{C}}$ the sufficiency of considering the differential equations

$$\delta'(\omega')\varphi = \lambda^2\varphi \quad (= \gamma_n(\omega')(\lambda) \cdot \varphi)$$

in our study of spherical functions on G may be justified as follows.

2.10 Theorem([2.], p. 145). If $\varphi_{\lambda}, \lambda \in \mathfrak{a}_{\mathbb{C}}^*$ is an eigenfunction of $\delta'(\omega')$, then it is also an eigenfunction of $\delta'(q), q \in \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, with the same eigenvalue.

Proof. We first show that the differential operators, $\delta'(q), q \in \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, commute with each other. Indeed, for any $q_1, q_2 \in \mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$, and any $f \in C^{\infty}(G^+//K)$, the commutativity of $\mathfrak{Q}(\mathfrak{g}_{\mathbb{C}})$ implies that $q_1 q_2 f = q_2 q_1 f$. Hence

$$\delta'(q_1 q_2) \tilde{f} = \delta'(q_2 q_1) \tilde{f},$$

where $\tilde{f} = f|_{A^+}$. Therefore

$$\delta'(q_1) \delta'(q_2) = \delta'(q_2) \delta'(q_1) \cdots (*)$$

Now let $\varphi_{\lambda}, \lambda \in \mathfrak{a}_{\mathbb{C}}^*$, be an eigenfunction of $\delta'(\omega')$. i.e., for some $\lambda \in \mathbb{C}$, we have $\delta'(\omega')\varphi_{\lambda} = \gamma(\omega')(\lambda) \cdot \varphi_{\lambda}$. Thus

$$\delta'(\omega')(\delta'(q)\varphi_{\lambda}) = \delta'(q)(\delta'(\omega')\varphi_{\lambda}) = \gamma(\omega')(\lambda)(\delta'(q)\varphi_{\lambda}),$$

meaning that $\delta'(q)\varphi_{\lambda}$ is an eigenfunction of $\delta'(\omega')$. However, by the uniqueness of Theorem 2.9, we must have that $\delta'(q)\varphi_{\lambda}$ is a constant multiple of φ_{λ} , as required. \square

The above result explains that it is sufficient to consider the operator

$$\delta'(\omega') = \tilde{\omega}' = \frac{d^2}{dt^2} + 2 \frac{\cosh 2t}{\sinh 2t} \frac{d}{dt} + 1$$

in the study of spherical functions, φ_{λ} , on $SL(2, \mathbb{R})$. In this case we have $\gamma(\omega')(\lambda) = \lambda^2$ so that the equation

$$\delta'(\omega')\varphi_{\lambda} = \gamma_n(\omega')(\lambda) \cdot \varphi_{\lambda}$$

becomes

$$\frac{d^2\varphi_\lambda}{dt^2} + 2\frac{\cosh 2t}{\sinh 2t}\frac{d\varphi_\lambda}{dt} + (1 - \lambda^2)\varphi_\lambda = 0 \cdots (\Upsilon)$$

i.e.,

$$\sinh^2(2t)\frac{d^2\varphi_\lambda}{dt^2} + 2\sinh 2t \cosh 2t\frac{d\varphi_\lambda}{dt} + \sinh^2(2t)(1 - \lambda^2)\varphi_\lambda = 0$$

Now setting $z = \cosh 2t$, and defining $\varphi_\lambda(t)$ as $\Phi_\lambda(z)$, we would have $\frac{d\varphi_\lambda}{dt} = (2\sinh 2t)\frac{d\Phi_\lambda}{dz}$ and $\frac{d^2\varphi_\lambda}{dt^2} = 4\sinh^2(2t)\frac{d^2\Phi_\lambda}{dz^2} + 4\cosh 2t\frac{d\Phi_\lambda}{dz}$ so that the last differential equation above transforms to

$$\sinh^4(2t)\frac{d^2\Phi_\lambda}{dz^2} + 2\sinh^2(2t)\cosh(2t)\frac{d\Phi_\lambda}{dz} + \frac{1}{4}\sinh^2(2t)(1 - \lambda^2)\Phi_\lambda = 0.$$

We finally have, with $\sinh^2(2t) = -(1 - z^2)$,

$$(1 - z^2)\frac{d^2\Phi_\lambda}{dz^2} - 2z\frac{d\Phi_\lambda}{dz} + \frac{(\lambda^2 - 1)}{4}\Phi_\lambda = 0.$$

This is the well-known *Legendre equation*.

This confirms that the spherical functions on $G = SL(2, \mathbb{R})$ are essentially the Legendre functions as enunciated in [4.], pp. 405 – 407.

It is a well-known fact in the general theory of ordinary differential equations that we can now consider the associated *confluent Legendre functions* on $G = SL(2, \mathbb{R})$ in this context, and since every spherical function on $G = SL(2, \mathbb{R})$ is a Legendre function, we may refer to the confluent Legendre functions on $G = SL(2, \mathbb{R})$ as *confluent spherical functions*. This is the motivation for the next section where this outlook is generated to all semisimple Lie groups with real rank 1.

§3. Reduction to the real rank 1 case.

It is appropriate, from Theorem 2.10, to find the general Casimir operator for the semisimple Lie group G in order to get the generalization of the Legendre equation of the group $G = SL(2, \mathbb{R})$ as already seen above. To this end let J be the two-sided proper ideal of $U(\mathfrak{g}_\mathbb{C})$, generated over \mathbb{C} by elements of the form $X \otimes Y - Y \otimes X$, where $X, Y \in \mathfrak{g}_\mathbb{C}$. The quotient $T(\mathfrak{g}_\mathbb{C})/J$ is the *symmetric algebra* of $\mathfrak{g}_\mathbb{C}$, denoted as $\mathcal{S}(\mathfrak{g}_\mathbb{C})$.

Clearly if $\mathfrak{g}_{\mathbb{C}}$ is abelian, then $U(\mathfrak{g}_{\mathbb{C}}) = \mathcal{S}(\mathfrak{g}_{\mathbb{C}})$. Even if $\mathfrak{g}_{\mathbb{C}}$ is non abelian, so that we only have $U(\mathfrak{g}_{\mathbb{C}}) \supset \mathcal{S}(\mathfrak{g}_{\mathbb{C}})$ in general, there is a map, $\lambda : \mathcal{S}(\mathfrak{g}_{\mathbb{C}}) \longrightarrow U(\mathfrak{g}_{\mathbb{C}})$, called the *Harish-Chandra symmetrization map* given as

$$\lambda(X_1 \cdots X_r) = \frac{1}{r!} \sum_{\sigma} X_{\sigma(1)} \cdots X_{\sigma(r)},$$

where $X_1, \cdots, X_r \in \mathfrak{g}_{\mathbb{C}}$ and σ runs over the set of all permutations of the set $\{1, \cdots, r\}$. This is a linear isomorphism and it may be shown that, for every $x \in G$, we have $\lambda \circ Ad(x) = Ad(x) \circ \lambda$, where $Ad(x)$ is viewed as a map from $\mathcal{S}(\mathfrak{g}_{\mathbb{C}})$ into $U(\mathfrak{g}_{\mathbb{C}})$ (By an adaptation of Theorem 2.2 to $\mathcal{S}(\mathfrak{g}_{\mathbb{C}})$)([4], p. 393). We then have the following.

3.1 Theorem([13.], p. 103). If $I(\mathfrak{g}_{\mathbb{C}})$ is the subset of members of $\mathcal{S}(\mathfrak{g}_{\mathbb{C}})$ which are $Ad(G)$ -invariant, then $\lambda : I(\mathfrak{g}_{\mathbb{C}}) \longrightarrow \mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ is a linear isomorphism.

Proof. This is a direct consequence of the relation $\lambda \circ Ad(x) = Ad(x) \circ \lambda$, for $x \in G$. \square

The above result allows us to use a basis of $I(\mathfrak{g}_{\mathbb{C}})$ in the construction of a basis of $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$. Since it is known that $\mathcal{S}(\mathfrak{g}_{\mathbb{C}})$ is essentially the polynomial algebra on $\mathfrak{g}_{\mathbb{C}}$, we may introduce the *Casimir polynomial*, ξ , on $\mathfrak{g}_{\mathbb{C}}$ given as $\xi(Z) = tr(adZ)^2$. It follows that $\xi \in \mathcal{S}(\mathfrak{g}_{\mathbb{C}})$ and hence we have $\lambda(\xi) \in \mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$. If we define $\tilde{\xi}$ on $\mathfrak{g}_{\mathbb{C}}$ by the requirement

$$\langle \tilde{\xi}, Z \rangle = \xi(Z), Z \in \mathfrak{g}_{\mathbb{C}},$$

then we shall refer $\lambda(\tilde{\xi})$ as the Casimir operator and denote it by ω . The situation for $\mathfrak{g} = sl(2, \mathbb{R})$ may be used to justify these terms. Indeed, the Casimir polynomial in this example is $H^2 + 4YX$ and, hence, $\lambda(H^2 + 4YX) = \lambda(HH) + 4\lambda(YX) = \frac{1}{2!}(HH + HH) + 4[\frac{1}{2!}(YX + XY)] = H^2 + 2YX + 2XY = H^2 + 2YX + (2H + 2YX) = H^2 + 2H + 4YX$, which when normalized gives exactly ω' . In the general case we have the following, where we denote B as the Cartan-Killing form on $\mathfrak{g}_{\mathbb{C}} \times \mathfrak{g}_{\mathbb{C}}$.

3.2 Theorem([12.], p. 217). The Casimir operator ω belongs to the center, $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ of $\mathfrak{U}(\mathfrak{g}_{\mathbb{C}})$. If $\{X_1, \cdots, X_m\}$ is a basis for $\mathfrak{g}_{\mathbb{C}}$ and $\{X^1, \cdots, X^m\}$ is the dual basis defined by $B(X_i, X^j) = \delta_{ij}$, then

$$\omega = \sum_{1 \leq i \leq m} X_i X^i.$$

Proof. The first statement holds from the linear isomorphism in Theorem 3.1. The dual basis, $\{X^1, \cdots, X^m\}$, exists since B is a non-singular symmetric bilinear form.

Now for every $X \in \mathfrak{g}_{\mathbb{C}}$ we have that $X = \sum_{1 \leq j \leq m} B(X, X^j) X_j$ so that

$$(adX)^2 = \sum_{1 \leq r, s \leq m} B(X, X^r) B(X, X^s) adX_r adX_s.$$

Therefore

$$\xi(X) = tr(adX)^2 = \sum_{1 \leq r, s \leq m} B(X, X^r) B(X, X^s) B(X_r, X_s).$$

Hence $\tilde{\xi} = \sum_{1 \leq r, s \leq m} B(X_r, X_s) X^r X^s$. We then have that $\omega = \lambda(\tilde{\xi})$
 $= \sum_{1 \leq r, s \leq m} B(X_r, X_s) X^r X^s = \sum_{1 \leq s \leq m} (\sum_{1 \leq r \leq m} B(X_r, X_s) X^r) X^s = \sum_{1 \leq s \leq m} X_s X^s$,
as expected. \square

We need to now compute the expression for the constant coefficient differential operator, $\gamma(\omega)$, for ω as in Theorem 3.2 above, in anticipation of its use in the formula for $\delta'(\omega)$ contained in Theorem 2.9. This would require using a specific basis of \mathfrak{g} according to the decomposition $\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{a} \oplus \sum_{\lambda \in \Delta} \mathfrak{g}_{\lambda}$, as we now proceed next.

Let $\{Z_i\}_{1 \leq i \leq m}$ be a basis of \mathfrak{m} such that $B(Z_i, Z_j) = -\delta_{ij}$, and let $\{H_i\}_{1 \leq i \leq a}$ be a basis of \mathfrak{a} such that $B(H_i, H_j) = \delta_{ij}$. This is possible since B is negative-definite on \mathfrak{t} (hence on \mathfrak{m}) and positive-definite on \mathfrak{p} (hence on \mathfrak{a}). Also since, for any two roots α, β of $(\mathfrak{g}, \mathfrak{a})$, the root-space \mathfrak{g}_{α} is orthogonal to \mathfrak{g}_{β} whenever $\alpha \neq \beta$, and since B is non-degenerate when restricted to $\mathfrak{g}_{\alpha} \times \mathfrak{g}_{-\alpha}$ (see [4.], p. 141) we may select a basis $\{X_{\alpha, i}\}_{1 \leq i \leq \dim(\mathfrak{g}_{\alpha})}$, of \mathfrak{g}_{α} , such that $B(X_{\alpha, i}, X_{-\alpha, i}) = \delta_{ij}$, where $\theta X_{\alpha, i} = X_{-\alpha, i}$ for a Cartan involution, θ , on \mathfrak{g} , for every root α . It is clear, from the assertion of Theorem 3.2, that $\omega|_{\mathfrak{m}} = \sum_{1 \leq i, j \leq m} Z_i Z_j = -(Z_1^2 + \dots + Z_m^2)$, $\omega|_{\mathfrak{a}} = \sum_{1 \leq i, j \leq a} H_i H_j = H_1^2 + \dots + H_a^2$ and $\omega|_{\mathfrak{g}_{\alpha}} = \sum_{1 \leq i, j \leq \dim(\mathfrak{g}_{\alpha})} X_{\alpha, i} X_{\alpha, j} = \sum_{1 \leq i, j \leq \dim(\mathfrak{g}_{\alpha})} (X_{\alpha, i} X_{-\alpha, i} + X_{-\alpha, i} X_{\alpha, i})$. Hence the direct sum $\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{a} \oplus \sum_{\lambda \in \Delta} \mathfrak{g}_{\lambda}$ now implies that

$$\omega = -(Z_1^2 + \dots + Z_m^2) + (H_1^2 + \dots + H_a^2) + \sum_{1 \leq i \leq n(\alpha), \alpha > 0} (X_{\alpha, i} X_{-\alpha, i} + X_{-\alpha, i} X_{\alpha, i}).$$

The present form of the Casimir operator in these bases may now be used to compute the constant coefficient differential operator, $\gamma(\omega)$.

3.3 Lemma ([2.], p. 94). In the above form of ω , we have that

$$\gamma(\omega) = \sum_{1 \leq i \leq a} H_i^2 - B(H_{\rho}, H_{\rho})$$

where $H_\rho = \frac{1}{2} \sum_{\alpha>0} n(\alpha)H_\alpha$ and H_α is uniquely defined by the requirement, $\alpha(H) = B(H_\alpha, H)$ for all $H \in \mathfrak{a}$.

Proof. We know that $X_{\alpha,i}X_{-\alpha,i} = X_{-\alpha,i}X_{\alpha,i} + [X_{\alpha,i}, X_{-\alpha,i}]$. Now if $X \in \mathfrak{g}_\alpha$ and $X' \in \mathfrak{g}_{-\alpha}$, then $[X, X'] \in \mathfrak{g}_{\alpha=0} \subset \mathfrak{m} \oplus \mathfrak{a}$, and, for every $H \in \mathfrak{a}$, we always have $B(H, [X, X']) = \alpha(H)B(X, X')$. These sum up to give $[X, X'] \equiv B(X, X')H_\alpha(\text{mod } \mathfrak{m})$, for every $X \in \mathfrak{g}_\alpha, X' \in \mathfrak{g}_{-\alpha}$. In particular

$$[X_{\alpha,i}, X_{-\alpha,i}] \equiv B(X_{\alpha,i}, X_{-\alpha,i})H_\alpha(\text{mod } \mathfrak{m}) = H_\alpha(\text{mod } \mathfrak{m}).$$

It then follows that

$$\begin{aligned} \omega &= -\sum_{k=1}^m Z_k^2 + \sum_{i=1}^a H_i^2 + \sum_{1 \leq i \leq n(\alpha), \alpha > 0} (X_{\alpha,i}X_{-\alpha,i} + X_{-\alpha,i}X_{\alpha,i}) \\ &= -\sum_{k=1}^m Z_k^2 + \sum_{i=1}^a H_i^2 + \sum_{1 \leq i \leq n(\alpha), \alpha > 0} (2X_{-\alpha,i}X_{\alpha,i} + [X_{\alpha,i}, X_{-\alpha,i}]) \\ &\equiv -\sum_{k=1}^m Z_k^2 + \sum_{i=1}^a H_i^2 + \sum_{1 \leq i \leq n(\alpha), \alpha > 0} (2X_{-\alpha,i}X_{\alpha,i} + H_\alpha(\text{mod } \mathfrak{m})). \end{aligned}$$

Now using the fact that $\mathfrak{Q}(\mathfrak{g}_\mathbb{C}) \cap (\mathfrak{t}U(\mathfrak{g}_\mathbb{C}))$, which contains \mathfrak{m} , is the kernel of $\beta_{\mathfrak{n}}$, we have

$$\beta_{\mathfrak{n}} = -0 + \sum_{i=1}^a H_i^2 + \sum_{\alpha>0} n(\alpha)H_\alpha.$$

Hence

$$\begin{aligned} \gamma(\omega)(\lambda) &= \beta_{\mathfrak{n}}(\omega)(\lambda - \rho) \\ &= \beta_{\mathfrak{n}}(\omega)(\lambda) - \beta_{\mathfrak{n}}(\omega)(\rho) \\ &= \left(\sum_{i=1}^a H_i^2 + \sum_{\alpha>0} n(\alpha)H_\alpha \right) - \left(2 \sum_{1 \leq i \leq a} \rho(H_i)H_i - \sum_{1 \leq i \leq a} \rho(H_i)^2 + \sum_{\alpha>0} n(\alpha)\rho(H_\alpha) \right). \end{aligned}$$

i.e.,

$$\gamma(\omega) = \sum_{i=1}^a H_i^2 + \sum_{\alpha>0} n(\alpha)H_\alpha - 2 \sum_{1 \leq i \leq a} \rho(H_i)H_i + \sum_{1 \leq i \leq a} \rho(H_i)^2 - \sum_{\alpha>0} n(\alpha)\rho(H_\alpha).$$

If we now observe, due to the orthonormality of the basis $\{H_i\}$ of \mathfrak{a} relative to B , that $\sum_{1 \leq i \leq a} \rho(H_i)H_i = H_\rho = \frac{1}{2} \sum_{\alpha > 0} n(\alpha)H_\alpha$ and $\sum_{1 \leq i \leq a} \rho(H_i)^2 = B(H_\rho, H_\rho) = \frac{1}{2} \sum_{\alpha > 0} n(\alpha)\rho(H_\alpha)$, we now have

$$\begin{aligned} \gamma(\omega) &= \sum_{i=1}^a H_i^2 + \sum_{\alpha > 0} n(\alpha)H_\alpha - 2\left(\frac{1}{2} \sum_{\alpha > 0} n(\alpha)H_\alpha\right) + \left(\frac{1}{2} \sum_{\alpha > 0} n(\alpha)\rho(H_\alpha)\right) - \sum_{\alpha > 0} n(\alpha)\rho(H_\alpha) \\ &= \sum_{i=1}^a H_i^2 - \frac{1}{2} \sum_{\alpha > 0} n(\alpha)\rho(H_\alpha) = \sum_{i=1}^a H_i^2 - B(H_\rho, H_\rho). \quad \square \end{aligned}$$

We can therefore get the expression for the radial component, $\delta'(\omega)$, of ω using the above expression for $\gamma(\omega)$ in Theorem 2.9. This would give the most general form of the formula

$$\delta'(\omega') = \frac{d^2}{dt^2} + 2 \frac{\cosh 2t}{\sinh 2t} \frac{d}{dt} + 1$$

of the case of $G = SL(2, \mathbb{R})$ to all connected semisimple Lie group G with finite center. The result is as follows.

3.4 Proposition ([2.], p. 133, and [3(a.)], p.269). The radial component, $\delta'(\omega)$, of the differential operator ω is given as

$$\delta'(\omega) = \sum_{1 \leq i \leq a} H_i^2 + 2 \sum_{\alpha > 0} n(\alpha)g_\alpha H_\alpha + \sum_{\alpha > 0} n(\alpha)H_\alpha.$$

Proof. Having known that

$$\omega = - \sum_{k=1}^m Z_k^2 + \sum_{i=1}^a H_i^2 + \sum_{1 \leq i \leq n(\alpha), \alpha > 0} (X_{\alpha,i}X_{-\alpha,i} + X_{-\alpha,i}X_{\alpha,i})$$

and that the first sum vanishes under β_n , it remains for us to find an expression for the sum $\sum_{1 \leq i \leq n(\alpha), \alpha > 0} (X_{\alpha,i}X_{-\alpha,i} + X_{-\alpha,i}X_{\alpha,i})$ in terms of members, f_α and g_α , of \mathcal{R}_0^+ , as expected in the expression for $\delta'(\omega)$ in Theorem 2.9 and explicitly seen in the case of $G = SL(2, \mathbb{R})$.

To this end, let $X_{\alpha,i} = K_{\alpha,i} + S_{\alpha,i}$ where $K_{\alpha,i} \in \mathfrak{t}$ and $S_{\alpha,i} \in \mathfrak{p}$. Since $[X_{\alpha,i}, X_{-\alpha,i}] = [K_{\alpha,i} + S_{\alpha,i}, K_{-\alpha,i} + S_{-\alpha,i}] = [K_{\alpha,i} + S_{\alpha,i}, -K_{\alpha,i} + S_{\alpha,i}] = 2[K_{\alpha,i}, S_{\alpha,i}] \in \mathfrak{p}$, and $[X, X'] \equiv B(X, X')H_\alpha \pmod{\mathfrak{m}}$, $X \in \mathfrak{g}_\alpha$, $X' \in \mathfrak{g}_{-\alpha}$, we arrive at $[X_{\alpha,i}, X_{-\alpha,i}] = H_\alpha$. This then means that $[K_{\alpha,i}, X_{\alpha,i}] = [K_{\alpha,i}, S_{\alpha,i}] = \frac{1}{2}[X_{\alpha,i}, X_{-\alpha,i}] = \frac{1}{2}H_\alpha$. Now setting $X = X_{\alpha,i}$ and $g = X_{-\alpha,i}$ in

$$\theta X \cdot g = f_\alpha(h)(X + \theta X)^{h^{-1}}g - g_\alpha(h)(X + \theta X)g$$

(from the equation (*) of the proof of Theorem 2.7) gives

$$\begin{aligned}
2X_{-\alpha,i}X_{\alpha,i} &= -2\theta(X_{\alpha,i})X_{\alpha,i} \\
&= -2[f_\alpha(h)(X_{\alpha,i} + \theta X_{\alpha,i})^{h^{-1}}X_{\alpha,i} - g_\alpha(h)(X_{\alpha,i} + \theta X_{\alpha,i})X_{\alpha,i}] \\
&= -2f_\alpha(h)[(K_{\alpha,i} + S_{\alpha,i}) + (K_{\alpha,i} - S_{\alpha,i})]^{h^{-1}}X_{\alpha,i} \\
&\quad + 2g_\alpha(h)[(K_{\alpha,i} + S_{\alpha,i}) + (K_{\alpha,i} - S_{\alpha,i})]X_{\alpha,i} \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)K_{\alpha,i}X_{\alpha,i} \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)(X_{\alpha,i}K_{\alpha,i} + [K_{\alpha,i}, X_{\alpha,i}]) \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)X_{\alpha,i}K_{\alpha,i} + 4g_\alpha(h)[K_{\alpha,i}, X_{\alpha,i}] \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)X_{\alpha,i}K_{\alpha,i} + 4g_\alpha(h)\left(\frac{1}{2}H_\alpha\right) \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)X_{\alpha,i}K_{\alpha,i} + 2g_\alpha(h)H_\alpha,
\end{aligned}$$

so that

$$\begin{aligned}
X_{\alpha,i}X_{-\alpha,i} + X_{-\alpha,i}X_{\alpha,i} \\
&= 2X_{-\alpha,i}X_{\alpha,i} + H_\alpha \\
&= -4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)X_{\alpha,i}K_{\alpha,i} + 2g_\alpha(h)H_\alpha + H_\alpha,
\end{aligned}$$

which when substituted into the above expression for ω gives

$$\omega = -\sum_{k=1}^m Z_k^2 + \sum_{i=1}^a H_i^2 + \sum_{1 \leq i \leq n(\alpha)} (-4f_\alpha(h)K_{\alpha,i}^{h^{-1}}X_{\alpha,i} + 4g_\alpha(h)X_{\alpha,i}K_{\alpha,i} + 2g_\alpha(h)H_\alpha + H_\alpha).$$

Using the kernel of β_n , and hence of γ_n , in the expression for $\delta'(\omega)$ we have

$$\begin{aligned}
\delta'(\omega) &= -0 + \sum_{i=1}^a H_i^2 - 4(0) + 4(0) + 2 \sum_{\alpha > 0} n(\alpha)g_\alpha(h)H_\alpha + \sum_{\alpha > 0} n(\alpha)H_\alpha \\
&= \sum_{i=1}^a H_i^2 + 2 \sum_{\alpha > 0} n(\alpha)g_\alpha(h)H_\alpha + \sum_{\alpha > 0} n(\alpha)H_\alpha. \quad \square
\end{aligned}$$

As from now on we may start the discussion on the asymptotic behaviour of the spherical functions, φ , from the perturbation theory of the system of differential equations satisfied by it given as

$$\delta'(\omega) \cdot \varphi_\lambda = \gamma_n(\lambda) \cdot \varphi_\lambda,$$

with $\delta'(\omega)$ as in Theorem 3.4. This is already contained in [3(a.)], [3(b.)], [11.] and, more recently [10.] In the present paper however, we shall seek to generalize the outlook that led, in §2., to the notion of a *confluent spherical function* to all semisimple Lie groups, with real rank 1. This would require loading the structure of G and its Lie algebra, \mathfrak{g} , into the known expressions for $\delta'(\omega)$ and $\gamma_n(\lambda)$ as we do next.

We now take G to be a connected semisimple Lie group with finite center and real rank 1. i.e., $\dim(\mathfrak{a}) = 1$. This implies that there exists exactly one simple root in Δ that we denote by α . This also means that 2α is the only other possible element in Δ^+ and, if p and q represents the numbers $n(\alpha)$ and $n(2\alpha)$ of roots in Δ which coincide on \mathfrak{a}^+ with α and 2α , respectively, then

$$\rho = \frac{1}{2} \sum_{\lambda \in \Delta^+} n(\lambda) \cdot \lambda = \frac{1}{2}(n(\alpha) \cdot \alpha + n(2\alpha) \cdot (2\alpha)) = \frac{1}{2}(p + 2q) \cdot \alpha$$

and that $p \geq 1, q \geq 0$. Now choose $H_0 \in \mathfrak{a}^+$ such that $\alpha(H_0) = 1$. Since, for any $H \in \mathfrak{a}$, we always have $B(H, H) = 2 \sum_{\lambda \in \Delta^+} n(\lambda) \cdot \lambda(H)^2 = 2(n(\alpha) \cdot (\alpha(H))^2 + n(2\alpha) \cdot ((2\alpha)(H))^2) = 2\alpha(H)^2(p + 4q)$, hence

$$B(H_0, H_0) = 2 \cdot \alpha(H_0)^2(p + 4q) = 2(p + 4q) \cdots \cdots (i.),$$

and $B(H_\alpha, H_\alpha) = 2 \cdot \alpha(H_\alpha)^2(p + 4q)$. i.e., $\alpha(H_\alpha) = 2\alpha(H_\alpha)^2(p + 4q)$. (since $\lambda(H') = B(H_\lambda, H')$, for all $H' \in \mathfrak{a}$.) This implies that $\alpha(H_\alpha) = (2(p + 4q))^{-1}$. Therefore

$$H_\alpha = \alpha(H_\alpha) \cdot H_0 = (2(p + 4q))^{-1} \cdot H_0 \cdots \cdots (ii.)$$

Equation (ii.) and the relation $\rho = \frac{1}{2}(p + 2q) \cdot \alpha$ also imply that $H_\rho = \rho(H_\alpha) \cdot H_0 = \frac{1}{2}(p + 2q) \cdot \alpha(H_\alpha) \cdot H_0 = \frac{1}{2}(p + 2q)(2(p + 4q))^{-1} \cdot H_0$. i.e.,

$$H_\rho = \frac{1}{2}(p + 2q)(2(p + 4q))^{-1} \cdot H_0 \cdots \cdots (iii.)$$

We now state the major result of this section as follows. This result generalizes the situation of $SL(2, \mathbb{R})$ above and motivates the concept of a confluent spherical function.

§4. Confluent Spherical Functions And Their Algebra.

4.1 Theorem. Let G be a real rank 1 connected semisimple Lie group with finite center and having the polar decomposition $G = K \cdot cl(A^+) \cdot K$. Then every K -biinvariant function on G is spherical iff it is a hypergeometric function.

Proof. Using the isomorphism $t \mapsto \exp tH_0, t \in \mathbb{R}$ between \mathbb{R} and A we identify H_0 with $\frac{d}{dt}$ so that

$$\begin{aligned} 2(p+4q)\delta'(\omega) &= 2(p+4q)[H_\alpha^2 + 2(n(\alpha)g_\alpha H_\alpha + n(2\alpha)g_{2\alpha}H_{2\alpha}) + (n(\alpha)H_\alpha + n(2\alpha)H_{2\alpha})] \\ &= 2(p+4q)[(2(p+4q))^{-1} \cdot H_0^2 + 2(p \cdot \xi_{-\alpha}(\xi_\alpha - \xi_{-\alpha})^{-1} \cdot (2(p+4q))^{-1} H_0 \\ &\quad + 2q \cdot \xi_{-2\alpha}(\xi_{2\alpha} - \xi_{-2\alpha})^{-1} (2(p+4q))^{-1} \cdot H_0) \\ &\quad + (p \cdot (2(p+4q))^{-1} \cdot H_0 + 2q \cdot (2(p+4q))^{-1} \cdot H_0)] \end{aligned}$$

(from an adaptation of formula (ii.) above)

$$= 2(p+4q)(2(p+4q))^{-1} \left[\frac{d^2}{dt^2} + 2(pg_1 \frac{d}{dt} + 2qg_2 \frac{d}{dt}) + (p \frac{d}{dt} + 2q \frac{d}{dt}) \right]$$

(where $g_k(t) = \xi_{-k\alpha}(t)(\xi_{k\alpha}(t) - \xi_{-k\alpha}(t))^{-1} = e^{-kt}(e^{kt} - e^{-kt})^{-1} = e^{-2kt}(1 - e^{-2kt})^{-1}, k = 1, 2$)

$$\begin{aligned} &= \frac{d^2}{dt^2} + \{2(pg_1 + 2qg_2) + (p + 2q)\} \frac{d}{dt} \\ &= \frac{d^2}{dt^2} + \{(2g_1 + 1)p + 2(2g_2 + 1)q\} \frac{d}{dt} = \frac{d^2}{dt^2} + \{p \coth t + 2q \coth 2t\} \frac{d}{dt} \\ &= \frac{d^2}{dt^2} + \{(p+q) \coth t + q \tanh t\} \frac{d}{dt} \dots \dots (iv) \end{aligned}$$

(using the relation $\coth 2t = \frac{1}{2}(\coth t + \tanh t)$)

In the case of $\gamma_n(\omega)(\lambda)$, we identify $\mathfrak{a}_\mathbb{C}^*$ with \mathbb{C} via the map $\lambda \mapsto \lambda(H_0)$, and for every $H_1 \in \mathfrak{a}$, we set $B(H_1, H_1) = 1$. Now if we substitute $H = H_1$ into the relation $B(H, H) = 2\alpha(H)^2(p+4q)$, it gives $H_1 = (2(p+4q))^{-\frac{1}{2}}H_0$, and when used with the

expression for H_ρ in (iii.) above we have (from Lemma 3.3) that

$$\begin{aligned}
\gamma_n(\omega)(\lambda) &= \lambda(H_1)^2 - B(H_\rho, H_\rho) \\
&= \lambda(H_1)^2 - \rho(H_\rho) \\
&= (\lambda((2(p+4q))^{-\frac{1}{2}}H_0))^2 - \rho\left(\frac{1}{2}(p+2q)(2(p+4q))^{-1}H_0\right) \\
&= ((2(p+4q))^{-1}\lambda(H_0))^2 - \left(\frac{1}{2}(p+2q)(2(p+4q))^{-1}\right)\rho(H_0) \\
&= (2(p+4q))^{-1}\lambda^2 - \left(\frac{1}{2}(p+2q)(2(p+4q))^{-1}\right)\left(\frac{1}{2}(p+2q)\alpha(H_0)\right) \\
&\quad \text{(from the expression } \rho = \frac{1}{2}(p+2q)\alpha)
\end{aligned}$$

$$= (2(p+4q))^{-1}\left(\lambda^2 - \frac{(p+2q)^2}{4}\right) \dots\dots (v.)$$

We now substitute the expressions for $\delta'(\omega)$ and $\gamma_n(\omega)(\lambda)$ in (iv.) and (v.) into the equation

$$\delta'(\omega)\varphi_\lambda = \gamma_n(\omega)(\lambda) \cdot \varphi_\lambda$$

and define the function f_λ on \mathbb{R} , as $f_\lambda(t) = \varphi_\lambda(\exp tH_0)$, which is possible because of the above isomorphism between \mathbb{R} and A , to have

$$\left(\frac{d^2}{dt^2} + \{(p+q)\coth t + q\tanh t\}\frac{d}{dt}\right)f_\lambda = \left(\lambda^2 - \left(\frac{(p+2q)^2}{4}\right)\right)f_\lambda \dots\dots (\Upsilon\Upsilon).$$

This is the equation (Υ) at the tail-end of §2. for $G = SL(2, \mathbb{R})$, where $p = 2$ and $q = 0$. We now transform ($\Upsilon\Upsilon$), as done in §2. for $\varphi_\lambda'' + 2\coth 2t\varphi_\lambda' + (1 - \lambda^2)\varphi_\lambda = 0$, by setting $z = -(\sinh t)^2$. This implies that $\frac{dz}{dt} = -2\sinh t \cosh t$ from which we may deduce that $\frac{d}{dt} = (-2\sinh t \cosh t)\frac{d}{dz}$ and $\frac{d^2}{dt^2} = (4\sinh^2 t \cosh^2 t)\frac{d^2}{dz^2} - 2(\sinh^2 t + \cosh^2 t)\frac{d}{dz}$. Defining a function, g_λ , on \mathbb{C} as $g_\lambda(z) = f_\lambda(t)$ under the transformation $z = -(\sinh t)^2$ then converts equation ($\Upsilon\Upsilon$) to

$$(z(z-1)\frac{d^2}{dz^2} + ((a+b+1)z-c)\frac{d}{dz} + ab)g_\lambda = 0,$$

where the constants a, b, c are given by $a = \frac{p+2q+2\lambda}{4}$, $b = \frac{p+2q-2\lambda}{4}$, $c = \frac{p+q+1}{2}$, respectively.

This is the well-known Gauss' hypergeometric equation. The point $z = 0$ which corresponds to $t = 0$, is a regular singular point for this equation, and it is known that

there is exactly one solution of it which is analytic at $z = 0$ and takes the value 1 there. This is the hypergeometric functions, $F(a, b, c : z)$, which, for $|z| < 1$, is given as

$$F(a, b, c : z) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k \cdot k!} z^k,$$

where $(m)_k := m(m+1) \cdots (m+k-1)$. Now since $g_\lambda(0) = f_\lambda(0) = \varphi_\lambda(1) = 1$ and g_λ is analytic in z at $z = 0$ we conclude that

$$\varphi_\lambda(\exp tH_0) = F(a, b, c : -(\sinh t)^2),$$

where $t \in \mathbb{R}$ and a, b, c are as given above.

Conversely, let a function φ_λ be K -biinvariant and be given as $\varphi_\lambda(\exp tH_0) = F(a, b, c : -(\sinh t)^2)$ for some a, b, c . If we consider the equation

$$\delta'(\omega)\varphi_\lambda = \gamma_n(\omega)(\lambda) \cdot \varphi_\lambda$$

for some yet-to-be known constant $\gamma_n(\omega)(\lambda) \in \mathbb{C}$, then $a = \frac{p+2q+2\sqrt{\gamma_n(\omega)(\lambda)}}{4}$, $b = \frac{p+2q-2\sqrt{\gamma_n(\omega)(\lambda)}}{4}$, and $c = \frac{p+q+1}{2}$. For any known real rank 1 connected semisimple Lie group G , with finite center, in which p and q are also known, we may solve for $\gamma_n(\omega)(\lambda)$ explicitly from the above relations. With the fact that $\varphi_\lambda(1) = F(a, b, c : 0) = 1$ we conclude that φ_λ is a spherical function on G . \square

The above result shows the one-to-one correspondence between the hypergeometric functions and spherical functions on real rank 1 semisimple Lie groups, G . Now we recall the well-known notion of the *confluent hypergeometric function* and use it, via Theorem 4.1 above, to introduce the notion of a *confluent spherical function* on G , which is then later generalized using the *Stanton-Tomas expansion* for spherical functions.

We recall that replacing $z (= -(\sinh t)^2)$ by $\frac{z}{b} (= \frac{-(\sinh t)^2}{b})$ the hypergeometric equation gives

$$z\left(z - \frac{z}{b}\right) \frac{d^2}{dz^2} g_\lambda(z) + \left\{c - \left(1 + \frac{a+1}{b}\right)z\right\} \frac{d}{dz} g_\lambda(z) - ag_\lambda(z) = 0$$

becoming, as $b \rightarrow \infty$,

$$z \frac{d^2}{dz^2} g_\lambda(z) + (c - z) \frac{d}{dz} g_\lambda(z) - ag_\lambda(z) = 0$$

whose solution, g_λ , is the confluent hypergeometric function, ${}_1F(a, c : z)$, is clearly given as

$${}_1F(a, c : z) = \lim_{b \rightarrow \infty} F(a, b, c : \frac{z}{b}) = \sum_{k=0}^{\infty} \frac{(a)_k}{(c)_k k!} z^k$$

where a, c and z are as above. Theorem 4.1 implies that there exists a K -biinvariant function, say φ_λ^σ , on G such that $\varphi_\lambda^\sigma = {}_1F$. It would be important to have a concise way of defining the function φ_λ^σ . To do this we study more closely the properties of ${}_1F$ as follows.

The relationship between z and t , given as $z = -(\sinh t)^2$, could be recast as $t = \sinh^{-1}(i\sqrt{z})$. Now the process of deriving the confluent hypergeometric equation above entails substituting z with $\frac{z}{b}$, before applying the limit as $b \rightarrow \infty$. Doing the same for the expression $t = \sinh^{-1}(i\sqrt{z})$, we have $t = \sinh^{-1}(i\sqrt{\frac{z}{b}})$. In applying the limit as $b \rightarrow \infty$, it follows that $t \rightarrow 0$. Now as $\lim_{b \rightarrow \infty} F(a, b, c : \frac{z}{b})$ gives ${}_1F(a, c : z)$, the last statement above implies that we study $\lim_{t \rightarrow 0} \varphi_\lambda(\exp tH_0)$. i.e., we study $\varphi_\lambda(\exp tH_0)$ for sufficiently small values of t . This observation is explicitly written as

$$\varphi_\lambda^\sigma(\exp tH_0) = \lim_{b \rightarrow \infty} \varphi_\lambda(\exp tH_0)|_{t=\sinh^{-1}(i\sqrt{\frac{z}{b}})}$$

Since $b \rightarrow \infty$ results to t being very small, the equality above implies that the study of the confluent spherical functions, φ_λ^σ , on semisimple Lie groups is the same as the study of spherical functions, φ_λ for sufficiently small values of t . i.e., the study of the function $\varphi_\lambda^\sigma : G \rightarrow \mathbb{C}$ in which given $\epsilon > 0$ we can find $\delta = \delta(\epsilon) > 0$ such that $|\varphi_\lambda(\exp tH_0) - \varphi_\lambda^\sigma(\exp tH_0)| < \epsilon$ whenever $t < \delta$.

This is the idea behind our notion of a confluent spherical function on G , and to develop this idea further we make use of the *Stanton-Tomas expansion* for spherical functions on a real rank 1 connected semisimple Lie group, with finite center (See [11.]). We however start with a motivation via the case of $G = SL(2, \mathbb{R})$ which proves the fact that *Legendre functions admit a series expansion in terms of Bessel functions*.

It has been shown by Harish-Chandra, [3(a.)], that every spherical function, on any connected semisimple Lie group G , with finite center and arbitrary real rank, has the integral expansion

$$\varphi_\lambda(x) = \int_K e^{(\lambda-\rho)(H(xk))} dk,$$

where $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$, $x \in G$, $\rho = \frac{1}{2} \sum_{\alpha \in \Delta^+} \dim(\mathfrak{g}_{\alpha}) \cdot \alpha$. When $G = SL(2, \mathbb{R})$, a calculation contained in [15.], p. 339, shows that

$$\varphi_{\lambda}(\exp tH_0) = \frac{1}{2\pi} \int_0^{2\pi} (\cosh t + \sinh t \cos \theta)^{(\lambda - \frac{1}{2})} d\theta.$$

This is the integral formula for the Legendre function, $P_{\lambda - \frac{1}{2}}(\cosh t)$ and is in consonance with the conclusion in §2. A change of contour in the integral yields

$$\varphi_{\lambda}(\exp tH_0) = c \int_0^t \cos(\lambda s) (\cosh t - \cosh s)^{-\frac{1}{2}} ds.$$

See [11.] for some details. Now for small values of t , it is known that

$$(\cosh t - \cosh s)^{-\frac{1}{2}} = (t^2 - s^2)^{-\frac{1}{2}} + \text{error}.$$

So that for sufficiently small values of t , we have

$$\varphi_{\lambda}(\exp tH_0) = c \int_0^t \cos(\lambda s) (t^2 - s^2)^{-\frac{1}{2}} ds = J_0(\lambda t),$$

where $J_n(\lambda t)$ is the Bessel function of order n , giving as the series expansion

$$J_n(x) = \frac{x^n}{2^n \cdot n!} \left\{ 1 - \frac{x^2}{2^2 \cdot 1!(n+1)} + \frac{x^4}{2^4 \cdot 2!(n+1)(n+2)} - \frac{x^6}{2^6 \cdot 3!(n+1)(n+2)(n+3)} + \dots \right\},$$

for any $n \in \mathbb{Z}$, with $J_{-n}(x) = (-1)^n J_n(x)$, and, if $n \notin \mathbb{Z}$,

$$J_n(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(n+k+1)} \left(\frac{x}{2}\right)^{n+2k}.$$

We can state our deductions above as follows.

4.2 Theorem. The confluent spherical functions, $\varphi_{\lambda}^{\sigma}$, on $G = SL(2, \mathbb{R})$ are the zero-th order Bessel functions, J_0 , of sufficiently small arguments.

Proof. Exactly as in the above deductions. \square

This idea generalizes to all real rank 1 semisimple Lie groups and is the first main result of Stanton and Tomas, [11.]. To state their result we make some preparations.

Let $n = \dim(G/K)$, which is known to be equal to $p + q + 1$, define $c_0 = c_0(G)$ and the function \mathcal{J}_{μ} , respectively, as

$$\pi^{\frac{1}{2}} 2^{(q/2)-2} \frac{\Gamma(\frac{n-1}{2})}{\Gamma(\frac{n}{2})}$$

and

$$\mathcal{J}_\mu(z) = \begin{cases} \frac{J_\mu(z)}{z^\mu} \Gamma(\mu + \frac{1}{2}) \Gamma(\frac{1}{2}) 2^{\mu-1}, & \text{if } z \neq 0, \\ 0, & \text{if } z = 0, \end{cases}$$

where J_μ is the Bessel function of order μ . Let also D be the Jacobian for the polar decomposition of G , then $D > 0$ on A^+ and is given by

$$D(t) = D(a_t) = e^{-\rho(\log a_t)} \cdot \prod_{\alpha > 0} (e^{2\alpha(\log a_t)} - 1)^{n(\alpha)}.$$

(see §2. for its reduction in the case of $SL(2, \mathbb{R})$)

In the case of a real rank 1 group, G , in which there are at most two positive roots, α and 2α , with multiplicity p and q , the Jacobian reduces to $D(t) = e^{-2\rho_0 t} g_1(t)^{-p} g_2(t)^{-q}$ where $g_k(t) = e^{-2kt}(1 - e^{-2kt})^{-1}$, $k = 1, 2$, and $\rho_0 = \frac{1}{2}(p + 2q)$. We now state a very important expansion formula for spherical functions, φ_λ , as follows.

4.3 Theorem([11.], p. 253) There exist $R_0 > 1, R_1 > 1$, such that for any t with, $0 \leq t \leq R_0$, the spherical function, φ_λ , has the given expansion

$$\varphi_\lambda(\exp tH_0) = c_0 \left[\frac{t^{n-1}}{D(t)} \right]^{1/2} \sum_{m=0}^{\infty} t^{2m} a_m(t) \mathcal{J}_{\frac{(n-2)}{2}+m}(\lambda t)$$

where

$$a_0(t) \equiv 1, \text{ and } |a_m(t)| \leq cR_1^{-m}. \quad \square$$

The error on truncating the above series is controlled as in the following.

4.4 Corollary([11.], p. 253) There exist $R_0 > 1, R_1 > 1$, such that for any t with $0 \leq t \leq R_0$ and any $M \geq 0$, the spherical function, φ_λ , is given as

$$\varphi_\lambda(\exp tH_0) = c_0 \left[\frac{t^{n-1}}{D(t)} \right]^{1/2} \sum_{m=0}^M t^{2m} a_m(t) \mathcal{J}_{\frac{(n-2)}{2}+m}(\lambda t) + E_{M+1}(\lambda t)$$

where

$$a_0(t) \equiv 1, \text{ and } |a_m(t)| \leq cR_1^{-m},$$

and

$$|E_{M+1}(\lambda t)| \leq c_M t^{2(M+1)},$$

if $|\lambda t| \leq 1$, and

$$|E_{M+1}(\lambda t)| \leq c_M t^{2(M+1)} \cdot (\lambda t)^{-\left(\frac{n-1}{2}\right)+(M+1)},$$

if $|\lambda t| > 1$. \square

We shall refer to the expansion in Theorem 4.3 above as the *Stanton-Tomas expansion* for spherical functions. We are therefore motivated to give the following general definition of a *confluent spherical function* on a real rank 1 semisimple Lie group G .

4.5 Definition. A *confluent spherical function* is any K -biinvariant function on $G(= K \cdot cl(A^+) \cdot K)$ which has the Stanton-Tomas expansion on A^+ .

Explicitly, a function $\varphi_\lambda^\sigma \in C(G//K)$ is a confluent spherical function on G if there exist $R_0 > 1$, $R_1 > 1$ such that for any t , with $0 \leq t \leq R_0$, and any $\lambda \in \mathbb{C}$,

$$\varphi_\lambda^\sigma(\exp tH_0) = c_0 \left[\frac{t^{n-1}}{D(t)} \right]^{1/2} \sum_{m=0}^{\infty} t^{2m} a_m(t) \mathcal{J}_{\frac{(n-2)}{2}+m}(|\lambda| t)$$

where $a_0(t) \equiv 1$, and $|a_m(t)| \leq cR_1^{-m}$.

We have $\varphi_0^\sigma(x) = 0$ while the introduction of complex λ 's and their moduli guarantee that $\varphi_{-\lambda}^\sigma(x) = \varphi_\lambda^\sigma(x)$. The differential equation satisfied by the confluent spherical function is contained in the following.

4.6 Theorem. The function, $g_\lambda : \mathbb{C} \rightarrow \mathbb{C}$ which coincides with φ_λ^σ on $K \times cl(A^+) \times K$, via the transformation $z = -(\sinh t)^2$, for sufficiently small values of t , is a solution of the differential equation

$$z \frac{d^2 y}{dz^2} + (c - z) \frac{dy}{dz} - ay = 0.$$

Proof. Since $z \frac{d^2}{dz^2} + (c - z) \frac{d}{dz}$ is a differential operator on \mathbb{C} we may take it as the realization of some $q \in \mathfrak{Q}$, under the transformation $z = -(\sinh t)^2$, for sufficiently small values of t . As φ_λ^σ is, in particular, a spherical function on G , with sufficiently small arguments, it satisfies the relation $q \cdot \varphi_\lambda^\sigma = \gamma(q)(\lambda) \varphi_\lambda^\sigma$, for some $\gamma(q)(\lambda) \in \mathbb{C}$. The result follows if we set $\gamma(q)(\lambda) = a$. \square

4.7 Remarks.

The confluent spherical functions, $\varphi_\lambda^\sigma(\exp tH_0)$, are defined for all $t \geq 0$, and all $\lambda \in \mathbb{C}$. Indeed, $\varphi_\lambda^\sigma(1) = \varphi_\lambda^\sigma(\exp(0)H_0) = 0$, since $D > 0$ on A^+ .

We denote the the set of all confluent spherical functions on G by $C^\sigma(G)$ and define the following operations on it:

- (i.) $\varphi_{\lambda_1}^\sigma + \varphi_{\lambda_2}^\sigma := \varphi_{\lambda_1 + \lambda_2}^\sigma$, for every $\varphi_{\lambda_1}^\sigma, \varphi_{\lambda_2}^\sigma \in C^\sigma(G)$.
- (ii.) $\alpha \varphi_\lambda^\sigma := \varphi_{\alpha\lambda}^\sigma$, for every $\varphi_\lambda^\sigma \in C^\sigma(G)$, $\alpha \in \mathbb{C}$.

(iii.) $\varphi_{\lambda_1}^\sigma \cdot \varphi_{\lambda_2}^\sigma := \varphi_{\lambda_1 \lambda_2}^\sigma$, for every $\varphi_{\lambda_1}^\sigma, \varphi_{\lambda_2}^\sigma \in C^\sigma(G)$.

One of our major results in this paper is the following. We denote the Schwartz algebra of spherical functions on G by $\mathcal{C}(G//K)$.

4.8 Theorem. The set $C^\sigma(G)$, endowed with the operations above, is an associative algebra over \mathbb{C} and there exists a non-trivial map $\mathcal{C}(G//K) \rightarrow C^\sigma(G)$.

Proof. We compute, using the operations above, that $C^\sigma(G)$ admits a \mathbb{C} -module and a ring structures, which are compatible. The non-trivial map is given as $\varphi_\lambda \rightarrow \varphi_\lambda^\sigma$. \square

We shall consider the extension of these results to a connected noncompact semisimple Lie group with finite center and of real rank $m > 0$ in another paper.

References.

- [1.] Barchini, L. and Zierau, R., *Differential operators on homogeneous spaces*, Notes of Lectures at the ICE-EM Australian Graduate School in Mathematics in Brisbane, 2nd-20th July, (2007). Retrieved from <http://www.math.okstate.edu/zierau/papers.html> on 12th January, 2011.
- [2.] Gangolli, R. and Varadarajan, V. S., *Harmonic analysis of spherical functions on real reductive groups*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, **101**, Springer-Verlag, 1988.
- [3.] Harish-Chandra, (a.) Spherical functions on a semisimple Lie group I, *Amer. J. Math.*, **80**, (1958), pp. 241 – 310. (b.) Spherical functions on a semisimple Lie group II, *Amer. J. Math.*, **80**, (1958), pp. 553 – 613.
- [4.] Helgason, S., *Differential geometry and symmetric spaces*, Academic Press, 1962.
- [5.] Hirano, M., Ishii, T., and Oda, T., confluent from Siegel-Whittaker functions to Whittaker functions on $Sp(2, \mathbb{R})$, (2004), pp. 1-14.

- [6.] Knapp, A. W., *Lie groups beyond an introduction*, Progress in Mathematics, **140**, Birkäuser Verlag, 2002.
- [7.] Knapp, A. W., and Trapa, P. E., *Representations of semisimple Lie groups*, IAS/Park City Mathematics Series, **8**, (1998), pp. 1-83.
- [8.] Lang, S., *$SL_2(\mathbb{R})$* , Addison-Wesley Publishing Company, 1975.
- [9.] Oyadare, O. O., Harmonic analysis of spherical functions on $SL(2, \mathbb{R})$, *Unpublished M.Sc. Dissertation*, University of Ibadan, 2007.
- [10.] Sawyer, P., The asymptotic expansion of spherical functions on symmetric cones, *Pacific J. Math.*, **200**(1), (2001), pp. 251-256.
- [11.] Stanton, R. J., and Tomas, P. A., Expansion for spherical functions on noncompact symmetric spaces, *Acta. Math.*, **140**, (1978), pp. 251-276.
- [12.] Varadarajan, V. S., *Lie groups, Lie algebras and their representations*, Springer-Verlag, 1984.
- [13.] Varadarajan, V. S., *An introduction to harmonic analysis on semisimple Lie groups*, Cambridge University Press, Cambridge, 1989.
- [14.] Warner, F., *Foundations of differentiable manifolds and Lie groups*, Scott, Foreman and Company, Glenview, 1971.
- [15.] Warner, G., *Harmonic analysis on semisimple Lie groups, II*, Grundlehren der mathematischen Wissenschaften, Band **189**, Springer-Verlag, 1972.
- [16.] Whittaker, E. T., and Watson, G. N., *A course of modern analysis*, 4 th edition, Cambridge University Press, Cambridge, 1965.