Unemployed persons' experiences and feedback of preventive health care services in Finland

Kirsi Niiranen¹, Hanna Hakulinen¹, Pirjo Manninen² and Kimmo Räsänen^{1,3}

Abstract

The aim of our study is to describe unemployed persons' experiences of preventive health care services, and to investigate how the health care system responds to their individual needs. The case is the Occupational Health Counselling project, Kuopio, Finland, in 2008-2010. The article is mainly representational. We examined from the perspective of the unemployed a transition one service to another, as well as the views of the unemployed on health and work ability. The link between health services and employment activities should be flexible between authorities. It seems that currently health check is an action still quite separate from employment activities. Client satisfaction with the service was high. Targeted at the unemployed, the special service type of health service meets clients' needs very well. Iindividuals who report reduced subjective health more attention should be offered. Young unemployed people especially benefit from objective health evaluations, and it imbues them self-confidence. Preventive health care services for the unemployed are important and services should be offered actively to unemployed. Finnish law requires municipalities to arrange these services but still they are not in every region. Also the process and contents of the services vary considerable. The situation has improved during the last few years, but work is still needed. Also, this study shows that preventive health care services for unemployed are of importance from the international point of view. The unemployed have many health care needs in common.

Key words

Unemployed, health services, network, client satisfaction

- ¹ Finnish Institute Of Occupational Health, OHS and Work Health, Kuopio, Finland
- ² MD practitioner, Kuopio Finland
- ^{1,3} University of Eastern Finland, School of Medicine, Kuopio, Finland

1. Introduction

The unemployed are a heterogeneous group and have different service needs [1]. All the problems of the unemployed cannot be solved by health care because the service structure is very complex and this group faces social problems [2]. Brussigemail et al (2014) showed that health promotion with labour market programmes is possible but requires agreements and coordination between the different social security branches. In labour market policy, stronger emphasis on voluntary programme participation is also needed. Health promotion requires skills and competencies; the staff in employment offices as well as in work organizations should adapt the needs of the unemployed [3]. Romppainen et al (2010) stated the production of health services for the unemployed to be a complex diversity of contemporary labor market trajectories. For example, non-standard employment and temporary work and unemployment are high, and some of the traditional professions have disappeared. In study occupational health nurses, felt that health care services are insufficient for unemployed clients of health services approaches. The study argues that although the structural and economic conditions can change relatively quickly, change in social habits is slow [4].

Attitudes in health care also have an impact. Welfare services do not always respond adequately to the needs of the unemployed [4]. Virtanen et al (2006) stated that the use of physician services varies according to labour market status, being relatively low among the non-permanently employed and the unemployed [5]. Dropping out of health services correlates strongly with unemployment according to Teff et al (2014) [6]. Policy makers should be aware of cyclical changes in preventive care use. As physician recommendations can have a strong impact on patients' use of health care, providers could increase efforts to persuade patients to seek screening examinations and necessary vaccinations during periods of high unemployment [6].

Reducing inequality in health services in Finland is one of the focus areas in development. Working-aged people's health care is divided into municipal health, occupational health and private health sector. Preventive services are used the more the higher the income group the person belongs to. However, persons in lower income groups and unemployed persons have more health problems. This relationship has been observed consistently across 23 European countries [7-8]. Many studies have demonstrated that differences between social classes in well-being and health have even grown [9-13]. Preventive health care services should be arranged according to Finnish Health Care Act [14] to unemployed in every region in Finland. Although about 89% of municipal health care centers carry out these services, the volume and quality of the practices are not known [15]. The coverage of occupational health care for salary and wage-earners in Finland in 2010 was 92% [16].

Researches related to the health of the unemployed and health services for them has been comprehensively studied but the studies on client's point of view are just a few [1, 3]. Aim of our study is to describe the unemployed persons' experiences of preventive health care services, and to investigate how the health care system responds to their individual needs. Case is the Occupational Health Counselling project Kuopio, Finland in 2008-2010. The article is mainly representational. It examines from the perspective of the unemployed a transition to one service to another, as well as the unemployed views on health and ability to work.

2. Materials and methods

2.1 The context of Occupational Health Counselling project

During 2007 – 2010, the unemployed health services for the unemployed were developed in 18 different sub-projects of the Finnish National Institute of Health and Welfare. One part of the

project was coordinated by Finnish Institute of Occupational Health called the Occupational Health Counselling [17]. This project took place in the years 2008 – 2010, and the preventive health services for the unemployed were organized in connection with the primary health care. Occupational Health Nurse (OHN) checked clients' health.

The role of the OHN is prominent and independent in Finland. They have the statutory education and qualification for required for this job. They coordinate the services and cooperate with organizations and enterprises. OHNs' are specialized in health promotion. [19]. They also deal with health examinations (over 600,000 every year) [20]. This means direct contact between the nurse and the employee. This usually involves a discussion of the employee's situation in life and the balance between private life and work life, workplace atmosphere, workload (e.g. time pressure, information load) and how the employee is coping with his/her work. The nurse also encourages the employee to adopt on safe work approaches and life habits and talks about nutrition, alcohol, tobacco and physical exercise. Working with unemployed needs know-how about unemployment and labour market` specific issues. [2, 17].

The OHN in health center made appointments with clients from various sectors, for example employment and economic development office, social welfare office, educational institutions and in the intermediate labor market (Figure 1). The clients were young unemployed people aged 15-25, and the middle-aged adults undergoing labour market training [2, 17, 18]. The intermediate labor market is middle face of a free labor market and unemployment. Unemployed persons aim to move to the free labor market. During the health examination the OHN surveyed clients' health and work ability by using different tests. After the meeting, the nurse and client planned in cooperation any further treatment or follow-up meetings, either in health care or other organizations (Figure 1) [2,18].




Figure 1. Occupational Health Counselling Modell

2.2 The data collection

Data was collected through interviews (n=9) with young clients, aged 15-25 and through questionnaire feedback; n=130 from all clients aged 15-54. We reminded once and also contacted

.

telephonically but still only 42 responded (response rate 32%). The majority were females (n = 37). The maximum responses was in 25-54 age groups. In the oldest age group were middle-aged adults undergoing labour market training in cleaning and assistant nursing. Those trades are dominated by women. The survey was conducted out in February-April 2010 (Annex 1).

A semi-structured interview was based on themes selected beforehand while there was no clear structure or order in the questions during the interview [21, 22]. This method of collecting material enables interaction between the interviewer and the interviewee, making it possible to use questions to extend understanding [23].

The interviews took 6-15 minutes. Three interviews were carried out early 2010 and four late 2010, and the remaining two in June 2011. Seven of the interviews were conducted out face to face and the remaining two by telephone (Annex 2).

We present our findings by combining the statistical analysis and content analysis. This triangulation method gives a wider understanding of the aim of the study. Also, the small number of responses of questionnaire does not allow more complex statistical analysis.

2.2.1 Content analysis

The interviews underwent content analysis using Atlas/ti program (Atlas/ti 5.1). Content analysis considers similarities and differences. [24, 25]. The units of analysis was chosen as the material purpose. In qualitative research with data collection, processing and analysis is difficult to separate [26, 27]. The process begins with the analysis of the analysis unit, which may be a word, a word combination or a phrase. The second step in the data is coding and the classification. In data classification simplified expressions are grouped, and the categories are classified as lower and higher. Consolidation will continue as long as it is meaningful. In the last phase of the material results are interpreted and conclusions are drawn. [26, 27, 28, 29].

The content analysis of the first phase of transcribed texts were fed into the Atlas/ti- program, after which the data were coded. The first phase of coding examined the content of the text and the meanings in the interviews. These parts were conceptualized with different codes, a total of 57. In the second part, the researcher examines whether codes have the correct code name. Some of the lower categories had the same contents, and the codes were combined. This was done manually by printing all of the coded expressions and by sorting, thus nine categories were established. As the data was small a two-level classification of the material was adequate.

2.2.2 Statistical analysis

SPSS was used for the statistical analysis. Data underwent descriptive analysis, direct distributions and the Pearson Exact Chi-Square test.

Statements based on a 5 step Likert scale (totally agree, tend to agree, neither agree nor disagree, tend to disagree, and strongly disagree). In the analysis we combined the first two classes. A comparison based through age distribution. The Pearson Exact Chi- Square test, value was used for statistical analysis because of the limited data.

3. Results

3.1 Experiences of service utility

The interviewed clients (n = 9) ranged from 20 to 25 years. Six of the clients were men. Their life situations ranged from unemployment (n=3), job training or education (n=4) to employment with the support of rehabilitative and work place units (n=2). Two of them had mental health problems or substance abuse problems. Some of the clients had moved to a new city, and they did not yet have a social network there. One client had no income at all at the moment, but he lived on his spouse's income. The common factor for all of them was rather low physical activity or lack of hobbies. Clients were directed to a health meeting from different sectors. The employment and Economic Development office sent three clients. Three other clients came from Intermediate Labor Market, and two clients had noticed a brochure of the project and had made an appointment by themselves. One client came suggested by his girlfriend.

Table 1. Background information of the clients who answered the follow-up questionnaire

	n=42		
Sex			
Female	37		
Male	5		
Age			
under 25	8		
25-34	4		
35-44	11		
45-54	19		
Life-situation			
unemployed	5		
employment	3		
with support			
employed	6		
education to	28		
employment			

By the time the clients were prompted their health checks, most of these have forgotten it. What they remembered, was fill in forms, measuring, height, weight and blood pressure and vaccination. Accessing to this service was easy and they all received basic information of their health status.

Of the clients 88%, opined that the health check gave a true picture of their health status. This was especially true for older clients. (Table 2).

All of the young clients interviewed felt that the service was useful and necessary. Most of these had rather limited suggestions for improving the service. Their opinions of health check meeting were mostly positive. They felt it important that they received an assurance that they had not more severe health problems. One client had concrete help in accessing treatment for back problems. "Really useful, you do not... Had there not been this opportunity for an appointment yet, just that.

[&]quot;It was at least quickly and the situation was uncomplicated and very clear. Easy to go there [laughs] "[sic] (Male 20).

And that it was free, that.... at their, there'd probably have to send us. It was fine to get that opportunity to go "[sic] (Male 21).

Clients were asked about the instructions during the health check. They considered instructions; lifestyle issues (smoking, alcohol, physical activity, and diet and weight management issues). Below are two comments about the instructions.

"Stop drinking. [laughing] I thought it was" [sic] (Male 23).

"Well, in this example from the physical activity of any kind that, you have a hobby activities, example from exercise according to all that the world of ball games and a lot of hobbies recommended that, in this age people can be recommended "[sic](Male 20).

Some of the interviewees felt that the health check also affected their health habits, although they were unable to specify this effects. One interviewee felt that the meeting import confidence and courage to apply for studies or work. Clients liked the easy way to get the service and the possibility to get their process forward. One client felt that health check was not relevant at all.

The feedback during the health check was considered sufficient by 73% of the respondents, most among the youngest age group, and least among the group 35-44 group (Table 2). Of the respondents, 40% considered that the health check had made them to pay more attention to health status and health habits. This was generally more common in the younger age groups (Table 2).

Table 2. Customer feedback of unemployed health check by age-group (n=42).

	Totally/ tend to agree					
	Age					
Opinion	Under 25	25–34	35–44	45–54	Total	p-value*
	n (%)	n (%)	n (%)	n (%)	n (%)**	
Health check gave a true picture of own health status	6 (75)	3 (75)	10 (90)	18 (95)	37 (88)	0,43
Feedback from the health check was sufficient	7 (88)	3 (75)	6 (55)	15 (79)	31 (73)	0,67
Health check made me to pay attention to the health status and health habits	4 (50)	2 (50)	4 (36)	7 (37)	14 (40)	0,93
Health check was necessary	7 (88)	3 (75)	6 (54)	17 (90)	33 (79)	0,24
Follow- up care had been achieved	4 (50)	2 (50)	6 (54)	9 (47)	21 (50)	1,00
Health examination attention was paid to the whishes	7 (88)	3 (75)	7 (64)	13 (68)	30 (71)	0,53
Health check should be regular	8 (100)	3 (75)	11 (100)	19 (100)	40 (98)	0,10
Unemployed need health checks	8 (100)	3 (75)	10 (91)	19 (100)	40 (95)	0,10
The results of health check were taken into	3 (38)	1 (25)	4 (36)	4 (21)	12 (29)	0,25
account in the employment services						

^{*} Pearson Exact Chi-Square test

The Beck Depression Inventory [30] showed over 10 points for six persons who responded to client feedback. Four of them totally agreed or tended to agree that the health check had prompted them to pay more focus to their health status and health habits.

^{**} All the respondents did not answer to all questions

The health check included regarding ability to work where compared to the life-time the best [30]. The scale was from 0 to 10. The estimation was under 8 (reasonable or poor) for four responses. Three of these agreed or tented to agree that the one outcome of the health check had made them to pay more attention to their health status and health habits.

Table 3. Client's opinion about the health check utility by age-group (n=42). Answer "Yes".

	Yes					
Health check was relevant	Age Under 25 n (%)	25–34 n (%)	35–44 n (%)	45–54 n (%)	Total n(%)**	p- value*
To my health	4 (50)	2 (50)	5 (45)	9 (47)	20 (48)	0,92
To my wellbeing	4 (50)	3 (75)	5 (50)	12 (67)	24 (60)	0,69
To my employment	2 (25)	1 (25)	3 (27)	4 (24)	10 (25)	0,98
To my finance	0 (0)	0 (0)	2 (18)	2 (12)	4 (10)	0,88

^{*} Pearson Exact Chi-Square test

Health check was considered necessary by 79% of the respondents. This was slightly more common among the youngest and the oldest age groups, but there was no statistical difference between the groups. (Table 2).

We asked whether the health check was meaningful (yes, no, cannot say) to their health, wellbeing, employment and finance. Naturally responses rated effects on wellbeing highest; 60%. Health (48%), employment (25%) and finance (10%) followed. (Table 3).

3.2 Follow up care and authority network

The respondents were asked about follow-up care, and there were no differences between the age groups (Table 2). Four of the six who had over 10 points in BDI considered the follow up care successful (agreed or tended to agree). Two out of four respondents had the score below 8 when estimating their ability to work (score 0-10). They tended to agree or totally agreed with follow-up care provision.

According clients' opinion, only 29% of the clients felt that the employment services tooking into account the results of the health check. This group tend to be more common among the youngest and the oldest age group. The interviewed young clients were not involved in multiactor meetings during the project. As a follow-up care included an appointment with the doctor or laboratory guidance.

The clients interviewed were asked about the network of authorities around them and other organizations in which their health or work ability issues had been discussed. Some of the young people did not have the need for network support, but listed a number of the authorities involved, such as the Social Insurance Institution (SII), the Employment and economic development office, the intermediate labor market, the social wealth ware services, the mental health services and doctors' appointments. Also, family members and clients' relatives were mentioned in connection with this theme.

^{**} All the respondents did not answer to all questions

"Okay, welfare, SII... Employment office, then." [sic] (Female, 22).

"SII and Social wealth ware they probably are, and, I'm not sure about vocational school network, it is anyway related"[sic] (Male 20).

3.3 The client role and wishes

Of the respondents, 71% considered their preferences were accommodated during the health examination. This was more common among the youngest age group (Table 2). In the older age group there were more expressions (32 %) of neither agree or disagree.

In the interviews, the client's own role rated as equal to the OHN. All of the interviewees were of the opinion that they gained relevant understanding of their life situation. One of them expressed a certain wonder regarding her situation in life. The OHN was described as a knowledgeable and attentive person.

Almost all of the clients conducted that a health check should be at regular intervals (98 %) and that unemployed persons need health checks (95%). Also, 86% considered that health checks were relevant to their own health; among the youngest age group, this percentage was smallest, 75 (Table 2).

However clients' opinion was that it is important this kind of service is also available for the unemployed and persons who are out of the coverage of occupational health services. The service was welcomed as it was. The only practical development proposal was that the number of questionnaires should be reduced.

"It's a good thing that there is a service, like this [sic] (Male 21).

In the free comments section of the feedback survey almost all clients hoped for regular health checks. One recommended that regular health checks should be mandatory. Clients hoped for more information about the services and health opportunities that unemployed could access to. One main positive point was that the service was free. Clients also hoped free exercise facilities and physiotherapist services to maintain their ability to work. Long appointment periods were criticized. One client felt that this kind of health check is representational of ones situation of life.

In the follow-up survey, 79% of the respondents thought that the health check was needed, 73% got sufficient feedback, and 88% considered that it gave a true picture of their health status; while 72% considered that their own needs and views were taken into account. Almost all considered that unemployed persons as a whole (95%) needed health checks. Almost all the respondents hoped regular health checks (Table 2).

4 Discussion

The aim of the study was to describe unemployed persons' experiences of the health care service and their opinions of maintaining work ability.

The unemployed need health checks, and they consider them important. 40% of the respondents thought that health check made them pay attention to their own health status and health habits. Of the six whose BDI was over 10 points, four of them agreed or tend to agree that the health check

had made them pay more attention to their own health status and health habits. Also, of the four whose estimation of work ability was under 8, three were this opinion. Meeting with a health professional is even more important and appropriate for these in a vulnerable life situation. Those persons need to get support and are willing to change their health habits. The young among unemployed specially benefit an objective health evaluation, and imports them self-confidence.

Axelsson et al, indicated that the quality of life (QOL) is good among a majority of unemployed young adults. Positive QOL also seems to be related to high self-esteem, satisfaction with spare time and a great extent of decision-making. In order to achieve more equity in health they suggest that special attention should be paid to individuals who report reduced subjective health, especially anxiety. Therefore, efforts should aim at empowering unemployed young adults by identifying their concerns and resources. Individual programs should include personal development beside work and education [31]. This has an important message also for health care also sector; personnel need to direct clients to right service.

According to client perceptions the results of the health check were taken into account by the employment services in only 29% of cases. Is an action still quite separate on the employment sector? The transition from the health check to the employment services is not necessary seamless even though the project tried to develop this co-operation. This link needs more attention in the future. Unemployed persons' health monitoring for maintaining work ability needs a complex networking of different actors from health care, social and employment sectors and voluntary participation. The health service should be close to the employment service, so that organizational limits disappear. The needs of the unemployed should be truly shared in different organizations. In this study we had a network of actors from various sectors. Referring organizations included, for example, the employment and economic development office, the social welfare office, educational institutions and the intermediate labor market. Based on the individual plan, the unemployed were referred to different actors in the health care sector, career counsellors, vocational psychologists, the Social Insurance Institution, the intermediate labor market among others.

The majority of the respondents thought that during the health examination attention was paid to their wishes. In the older age group, there were more expressions (32%) of neither agree nor disagree. Why is the older group more crucial than the younger group? This is perhaps because they have more life experience and they want more from the service. Young people find it difficult to compare services due to their lack of experience. Memory of the health checks was minimal. The health check should be more relevant and focused on clients' needs. A small group (4 persons) felt that the follow up care was not as successful as planned during the health check. New methods like mobile applications could prove useful even though they do not compensate for human contact. Even though some of health care personnel think that the health care of unemployed is mentally laborious these services should be available. It is also important to take care of the personnel's own coping, for example, by supervision of their work. Health care services should be offered actively to unemployed persons.

The report of Finnish National Institute of Health and Welfare report from regarding the health of those unemployed for a long time (Development Partnership Project 2007-2010) [1] showed the clients' feedback was very positive as regards their healths in the review. 61% (in this study 40%) of clients told in the review they paid more attention to the health status and health habits after the heath check. Client's transitions after a health often bound for the care and while some clients were moved to rehabilitation some others started receiving disability pension. In the partnership project, the clients were mainly long-term unemployed and belonged to the older age group [1]. Overall the results in this report and the current study are quite consistent.

The evidence from health checks has been recently discussed very much in the Occupational Health Service (OHS) field in Finland. There is scientific evidence about verified the effectiveness of health checks in OHS according to Taimela et al (2008, 2008, and 2010). They divided high`risk patients into an intervention group. After an one year follow-up, in the intervention group there were less sick leaves, and less costs than in control group [32, 33, and 34].

The benefits of health services for the unemployed has also been studied. For example, Kreuzfeld et al. (2012) [35] mentioned degrees in the blood pressure, CVD risk, back pain and depressive symptoms among long-term unemployed as a result physical activity program. Horns et al. (2012) [36] showed similar results, the healthy lifestyle after intervention. Rothländer et al. (2012) [37] used an experimental design, in which the intervention were used cognitive-behaviorism methods for health promotion. Researchers found a significant difference in the health status: the intervention group's self-rated health was better. Evidence can be found of the benefits of support of methods for the employment in Dalen et al. (2008) [38]. The similar evidence of workplace skill improvement in the unemployed in Finland has also been shown (Vuori et al., 2002) [39]. Based on these studies, participatory interventions seem to bring health and wellbeing benefits. Our study shows also welfare and health benefits for such clients.

This study and data has certain limitations. The response rate for the follow-up questionnaire was quite low (32%). The older age group answered more often than the younger group. It was hard to get responses even by phone. The young clients did not necessarily answer to a phone call or back after call request. The majority of respondents were women. The clients who were undergoing employment offices labour market training were assistant nurses or of those training to be cleaners. These lines of work are dominated by women. Because of the limited of the data, we could do only a simple statistical analysis. We could not compare, eg. differences between the genders in their opinion of the services. These results are approximate and cannot be generalized.

It was a challenge to get the participants for the interview. Although the interview time was agreed upon, the interviewee did not necessarily arrive at the venue. Reaching the interviewees by telephone was also difficult. The result of interviews were also rather limited and the interviewees responded to most items pretty briefly. The young clients did not have very many opinions about the service, and they were satisfied. This clientele is passive and do not necessarily count on authorities, not to mention researchers. In the content analysis data classification and naming was checked several times. Material classification, however, always represents researcher choices and interpretation [25]. When the interview data is relatively small, a two-level classification is sufficient. Also, direct quotes illustrate the material.

Verbeek et al (2001) [40] reviewed the literature on the concept of consumer satisfaction with health care and on surveys of satisfaction with occupational health care. Consumer satisfaction is an important indicator of quality in OHS. Researchers claim that high average ratings of satisfaction is the most important methodological problem to avoid. It is important to measure specific aspects of occupational healthcare experiences among patients and clients because satisfaction is a multidimensional concept. Client feedback helps to better understand client preferences and to improve the quality of the services. It would be important to inform these unemployed persons themselves of the study results.

Hall and Doman (1988) [41] have analyzed over 200 studies concerning patient/client satisfaction. The average "satisfaction" count was 76 %. Several researchers have showed that even though the satisfaction is 90 % the responses to certain specific questions are very critical and negative made in study [42, 43]. In this study we found also criticism, for example, in follow-up care.

5 Conclusion

Client satisfaction in health service for the unemployed was high. Targeted at the unemployed, this service meets clients' needs very well. Preventive health care services for the unemployed are important, and services should be offered actively. Individuals who report reduced subjective health should be given more attention. Young unemployed people especially benefit from objective health evaluations, and it imbues them self-confidence. The link between health care services and employment activities should be flexible between authorities.

Monitoring the health of unemployed persons in order to maintain their ability to work needs complex networking of different actors from health care, social and employment sectors and voluntary participation. It seems that currently health check is an action still quite separate from employment activities; the transition from health check is not necessary the seamless service with the employment service. This link needs more attention in the future. Voluntary program participation is also needed. This service construction is complex and needs new models in different organizations.

The findings give important information about preventive health care for the unemployed. In Finland, the law that stipulates municipalities arrange services, but certain regions still lack these. Also the process and contents of the services vary a lot. The situation has come better during the last few years, but work is still needed. From the international point of view this study shows that preventive health care services to unemployed should be arranged. The unemployed have many health care needs in common.

References

- [1] P.Saikku, "Terveyspalvelu työllistymisen tukena. Pitkäaikaistyöttömien terveystarkastukset- ja palvelut siirtymätyömarkkinoilla," Terveyden ja hyvinvoinnin laitos, Raportti, Helsinki. 22, 2009.
- [2] K. Niiranen, H. Hakulinen, P. Manninen and K. Räsänen, "Työttömien terveyspalvelujen kehittäminen verkosto mahdollisuutena," Työelämän tutkimus 2014, vol 12 no. 1 pp. 3-22.
- [3] M. Brussigemail, N.Draganoemail and S. Mümkenemail "Health promotion for unemployed jobseekers: New developments in Germany," Health Policy. 2014, vol. 114, no. 2-3, pp. 192–199.
- [4] K. Romppainen, A.Saloniemi, R.Jähi and P.Virtanen, "My health and treirs: clients constructing meanings for a health service programme for unemployed people," Sociology of Health & Illness. 2012 vol. 34 no. 6, pp. 809-825.
- [5] P. Virtanen, M. Kivimäki, J. Vahtera and M. Koskenvuo, "Employment status and differences in the one-year coverage of physician visits: different needs or unequal access to services?," BMC Health Serv Res. 2006, vol 2, no. 6, pp. 123.
- [6] N. Tefft, and A.Kageleiry, "State-Level Unemployment and the Utilization of Preventive Medical Services," Health Services Research Volume 2014, 49, no. 1, pp. 186–205.
- [7] C. Bambra and T.A. Eikemo, "Welfare state regimes, unemployment and health: a comparative study of the relationship between unemployment and self-reported health in 23 European countries," J Epidemiol Community Health 2009, 63:92-98.
- [8] J. Brown, E. Demou, M.A. Tristam, H. Gilmour, K. Sanati and E. Macdonald, "Employment Status and Health: Understanding the Health of the Economnically Inactive Population in Scotland", BMC Public Health 2012. 12, 327.
- [9] Health inequalities in EU [http://www.health-inequalities.eu/HEALTHEQUITY/EN/about_hi/health_inequalities/in_the_eu/ [acces 30.7.2014]
- [10] A. Kunst, EV. Bosand J.P. Mackenbach, J. P., "EU Working Group on Socio-economic Inequalities in Health. Monitoring socio-economic inequalities in health in the European Union: guidelines and illustrations. A report for the Health Monitoring Program of the European Commission," Department of Public Health, Erasmus University. Rotterdam, 2011.
- [11] T.Martelin, S. Koskinen and E. Lahelma, "Väestöryhmien väliset terveyserot. Suomalaisten terveys," 2005. 18.7.
- [12] T. Valkonen and T.M. Kauppinen, "Miesten kuolleisuuden alue-erot ja sosiaalinen segregaatio pääkaupunkiseudulla," Kvartti 2001, 1, pp. 7–21.
- [13] T. Heponiemi, M. Wahlström, M. Elovainio, T. Sinervo, A-M. Aalto and I. Keskimäki, "Katsaus työttömyyden ja terveyden välisiin yhteyksiin," Työ ja elinkeinoministeriön julkaisuja. Työ ja yrittäjyys 14, 2008
- [14] Terveydenhuoltolaki. Suomen säädöskokoelma 1326/2010.
- [15] L. Sinervo and L. Hietapakka, "Työttömien terveyspalvelut. Valtakunnallisen terveyskeskuskyselyn 2013 tulokset," Paper. Terveyden ja hyvinvoinnin laitos, 2013.

- [16] T. Kauppinen, P. Mattila-Holappa, M. Perkiö-Mäkelä, A. Saalo, J. Toikkanen, S. Tuomivaara, S. Uuksulainen, M. Viluksela and S. Virtanen, "Työ ja terveys Suomessa 2012," Työterveyslaitos. Tampere. Tammerprint Oy.2013.
- [17] L. Sinervo, "Työttömien terveyspalvelun juurrutus: Valtakunnallisen PTT-hankkeen kokemuksia, arviointitutkimuksen tuloksia ja kansallisia suosituksia," Terveyden ja hyvinvoinnin laitos, 2009.
- [18] K. Niiranen, H. Hakulinen, M.S. Huuskonen, A. Jahkola and K. Räsänen, "Supporting the work ability of the unemployed in the authority network," Journal of Social Medicine. Sosiaalilääketieteellinen aikakauslehti. 2008, 48, pp. 38–52.
- [19] Finnish Assossiation of Occupational Health Nurses http://www.stthl.net/fin/jasenyys/toimenkuva/ [acces 23.10.2014]
- [20] Kela työterveyshuoltotilasto. http://www.kela.fi/vuositilastot_kelan-tyoterveyshuoltotilasto. [acces 23.10.2014]
- [21]DF. Polit and BP. Hungler, "Nursing research Principles and Methods," London: J.B. Lippincott Company; 1987.
- [22] F. Fontana, "The Interview. From Structured Question to Negotiated Text," In Handbook of Qualitative Research. Edited by Edited by Denzin NK, Lincoln YS. Thousand Oaks: Sage Publications. 2000, pp. 645-672.
- [23] F. Marton, "Phenomenography describing conceptions of the world around us," Instructional Science 1981, 1981, 10, pp.177-200.
- [24] E. Latvala and L. Vanhanen-Nuutinen, "Laadullisen hoitotieteellisen tutkimuksen perusprosessi: sisällönanalyysi," Teoksessa Laadulliset tutkimusmenetelmät hoitotieteessä. Janhonen, S. & Nikkonen, M. (ed.) Juva. WS Bookwell Oy, 2003.
- [25] K. Krippendorff, "Content analysis. An Introduction to Its Methodology," Sage Publication, 1985.
- [26] J. Tuomi, J. and A. Sarajärvi, "Laadullinen tutkimus ja sisällönanalyysi," Jyväskylä. Gummerus Kirjapaino Oy, 2002.
- [27] M. I. Franklin, "Understanding Research: Coping with the Quantitative-Qualitative Divide," London/New York. Routledge, 2012.
- [28] J. Kylmä and T. Juvakka, "Laadullinen terveystutkimus," Edita Publishing Oy, 2012.
- [29] H. Kyngäs and L. Vanhanen, L, "Sisällön analyysi," Hoitotiede, 1999, 11, pp. 3–12.
- [30] Toimia- database http://www.thl.fi/toimia/tietokanta/mittari/tulokset/8 [acces 1.10.2014]
- [31] L. Axelsson and G. Ejlertsson, "Self-reported health, self-esteem and social support among unemployed young people a population-based study," Int J Soc Welfare 2002, vol. 11, no. 2, pp. 111-119.

- [32] S. Taimela, A. Malmivaara, S. Justén, E. Läära, H. Sintonen, J. Tiesko and T. Aro, "The effectiveness of two occupational health intervention programmes of two occupational health intervention programmes in reducing absence among employees at risk. Two randomized controlled trials," Occup Environm Med. 2008, vol. 65, no. 4, pp. 236–41
- [33] S. Taimela, S. Justén, P. Aronen, H. Sintonen, E. Läära, A. Malmivaara, J. Tiesko and T. Aro, "An occupational health intervention programme for workers at high risk for sicness absence. Cost effectiveness analysis based on randomized controlled trial," Occup Environm Med. 2008, vol. 65, no. 4, pp. 242-8.
- [34] S.Taimela, P. Aronen, A. Malmivaara, H. Sintonen, J. Tiesko and T. Aro T, "Effectiveness of a targeted occupational health intervention in workwers with high risk of sickness absence: baseline characteristics and adherence as effect modifying factors in a randomized controlled trial," J Occup Rehabil. 2010, vol. 20, no. 1, pp. 14-20.
- [35] Kreuzfeld, S., Preuss, M., Weippert, M. & Stoll, R, "Health effects and acceptance of a physical activity program for older long-term unemployed workers," International Archives of Occupational and Environmental Health 2012, vol. 85, no. 1, pp. 1-7
- [36] M.K. Horns, S.K. Seeger, M. Heinmüller, H. Limm, H.P. Waldhoff,, R. Salman, H. Gündel and P. Angerer, "Health promotion for long-term unemployed. Effects on motivation for a healthy lifestyle," Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2012, vol. 55, no. 5, pp. 728–738.
- [37] K. Rothländer, S. Mühlpfordt and P. Richter, "Evaluation of the health promotion program Active Coping with Unemployment (AktivA)," Zeitschrift für Gesundheitspsychologie 2012, vol. 20, no. 3, pp. 115–127.
- [38] R. Dale, B.K. Graversen and J.C.Van Ours, "Activating unemployed workers works; Experimental evidence from Denmark," Economics letters. Evaluation Frameworks for Development Programmes and Projects. 2008, vol. 100, no. 2, pp. 208-310.
- [39] J. Vuori, J. Silvonen, A.D.Vinokur, and R.P. Price, "The Työhön Job Search Program in Finland: Benefits for the unemployed with risk of depression or discournagement?," Journal of Occupational Health Psychology 2002b, 7, pp. 5–19.
- [40] J. Verbeek, van F. Dijk, K. Räsänen, H. Piirainen, E. Kankaanpää and C. Hulshof, "Consumer satisfaction with occupational health services: should it be measured?," OccupEnvironMed 2001, 58, pp. 272–278
- [41] J.A. Hall and M.C. Dornan, "Meta-analysis on satisfaction with nedical care: Description of research domain and analysis of overall satisfaction levels," Social Science Medicine 27. 1988, pp. 637-644.
- [42] S. Williams and M. Calnan, "Key determinants of consumer satisfaction with general practice," Fam. Practice 8, 1991.
- [43] R. Fitzpatrik, "Surveys of patient satisfaction. Important general considerations," British Medical Journal, 1991, vol. 302, 13.

ANNEX 1

The themes of follow-up questionnaire

- 1. Background information (sex, age, working situation)
- 2. Statements of the health check (Scale 1-5)
 - Health check gave a true picture of own health status
 - The practice worked well
 - Feedback from the health check was sufficient
 - Health check made me to pay attention to the health status and health habits
 - Health check was necessary
 - Follow- up care had been achieved
 - Health examination attention was paid to the whishes
 - Health check should be regular
 - Unemployed need health checks
 - The results of health check were taken into account in the employment services
- 3. Meaning of the health check (1= yes 2= no 3= I don't know)
 - To my health
 - To my wellbeing
 - To my employment
 - To my finance
- 4. Follow-up care after the health check (different follow-up services)
 - Physician
 - Health/sick care services
 - Dentist
 - Physiotherapist
 - Physical exercise services
 - Mental health clinic
 - Druck Abusers services
 - Labour Force services
 - Social wealth ware services
 - ISS services
 - Employment and Economic Development Office
 - Something else
- 5. Any other free comments

ANNEX 2

Theme interview questions to Occupational Health Counselling clients

- 1) Can you tell briefly about your life situation?
- 2) How did you come to Occupational Health Counselling project?
- 3) Can you tell in your own words what happened in the health meeting?
- 4) What good sides were there in the health meeting? What improvements would you suggest?
- 5) What kind of further treatment options were discussed?
- 6) How useful did you find the counselling?
- 7) What specific steps did you get from health appointment?
- 8) What kind of role/position did you have in health meeting? How was the understanding of your matters?
- 9) What other organizations have dealt with your health / your work ability related things?
- What kind of network operators are there around you?
- 10) How the health meeting has influenced with your health / your ability to work?
- 11) What new opportunities the health meeting highlighted to your health and work ability in the future?
- 12) What other feedback / wishes or development ideas you have about this service?