

# Modeling heteroscedastic, skewed and leptokurtic returns in discrete time

Ivivi Joseph Mwaniki

School of Mathematics, Statistics, Actuarial and Finance Division,  
University of Nairobi, Kenya

*Email: jimwaniki@uonbi.ac.ke*

August 3, 2015

## Abstract

Popular models of finance, fall short of accounting for most empirically found stylized features of financial time series data, such as volatility clustering, skewness and leptokurtic nature of log returns. In this study we propose a general framework for modeling asset returns which account for serial dependencies in higher moments and leptokurtic nature of scaled GARCH filtered residuals. Such residuals are calibrated to normal inverse Gaussian and hyperbolic distribution. Dynamics of risky assets assumed in Black Scholes model, Duans GARCH model and other benchmark models for contract valuation, are shown to be nested in the the proposed framework.

## Keywords

**Stylized facts, Normal inverse Gaussian, GARCH model, Hyperbolic distribution.**

## 1 Introduction

Uncertainty is central to much of modern finance theory. In option pricing for example, the uncertainty associated with future return of the underlying asset, is the most important determinant in the pricing function. Popular models such as the Black and Scholes [1973] model, based on the geometric Brownian motion have very nice mathematical properties which have been extensively used to value contracts. Empirical evidence suggest that the dynamics of the underlying process under the physical measure  $\mathbb{P}$  follow a more complicated process than the standard geometric Brownian motion with constant volatility.

Various studies have shown that the normal distribution does not accurately describe observed stock return data. In literature, for example Eberlein and Keller [1995], Barndorff-Nielsen [1998], Carr et al. [2002] and references therein, it is proposed that daily log returns, could be modeled by an exponential Lévy processes and geometric Lévy process, generating a lot of literature applied in pricing derivatives. Moreover, other studies suggest that a more improved model would include both stochastic volatility and a jump component (see Carr and Wu [2004]).

Over the past several decades, several stylized facts have emerged about the statistical behavior of speculative market daily returns such as aggregational Gaussianity, volatility clustering, changing variance, Taylor effect, leptokurtic residuals etc, see Rydberg [2000], Cont [2001], Tsay [2002], Stentoft [2008] and Rogers and Zhang [2011] for further documentation.

A typical finding concerning the return characteristic is that one period asset return, conditional on the most up-to-date information, exhibit a fat tail behaviour in addition to varying second order moments. The ARCH family of models introduced by Engle [1982] and generalized by Bollerslev [1986] has in the recent years gained prominence for modeling such dynamics. In the last few years, much interest has been given to the discrete-time GARCH option pricing models for instance see Pagan and Schwert [1990], Glosten et al. [1993], and Bollerslev et al. [1994]. Option pricing in GARCH models has been typically done using the local risk neutral valuation relationship (LRNVR) pioneered by Duan [1995]. The crucial assumptions in his GARCH model construction, are the conditional normal distribution of the asset returns under the physical probability measure  $\mathbb{P}$  and the invariance of the conditional volatility to the change of measure. The main objective of this study is to propose a general framework for modeling the underlying uncertainty driving heteroscedastic and leptokurtic daily stock market returns.

This article is organized as follows, the next section presents ARCH type model and class of generalized hyperbolic distribution for modeling some of the basic stylized facts of returns. In section three, we state the proposed general framework followed by several examples of popular benchmark models nested within the framework. Empirical analysis and parameter estimation are stated in section four, and section five concludes the study.

## 2 Preliminary considerations

Stock returns tend to exhibit a significant serial dependency in second moments. ARCH class models and several class of statistical distributions say NIG etc are known to models skewness and excess kurtosis of log returns.

### 2.1 ARCH models: Changing variance

ARCH model models have become popular for modeling financial time series because they are able to account for several empirical features like volatility clustering and leptokurtic in the distribution of returns. Most ARCH-type models involve a sequence of innovations whose variance is random. Conditioned on the past the variance depends only on the previous innovations and the previous conditional variances, and standard Wiener process generate the filtration. Most studies of daily stock returns using GARCH models and conditioned on normal distribution for the innovations; the re-scaled residuals showed excess kurtosis which violates the normality assumption. We investigate AR(1)-GARCH(1,1) model conditioned on normal distribution. Thus

$$\begin{aligned} X_t &= \log_e S_t - \log_e S_{t-1} \\ &= \hat{\mu} + \phi X_{t-1} + \sigma_t(Z_t + \mathfrak{L}_t), Z_t \sim N(0, 1) \\ \sigma_t^2 &= \omega + a\sigma_{t-1}^2 Z_{t-1}^2 + b\sigma_{t-1}^2 \end{aligned}$$

Parameter estimates  $\hat{\mu}, \phi, \omega, a, b$  are presented in Table 1 and Table 2.

A typical finding concerning the return characteristic is that one period asset return conditional on the most up to date information, continues to exhibit a fat tailed behaviour. This fat tailed behaviour is also known as conditionally leptokurtic, and can be modeled by a limiting class of generalized hyperbolic distribution.

## 2.2 Generalized hyperbolic distribution

The probability density function of the one-dimensional Generalized Hyperbolic distribution is given by the following:

$$f_{GH}(x; \alpha, \beta, \delta, \mu, \lambda) = \frac{(\gamma/\delta)^\lambda}{\sqrt{2\pi}K_\lambda(\delta\gamma)} \cdot \frac{K_{\lambda-\frac{1}{2}}(\alpha\sqrt{\delta^2 + (x-\mu)^2})}{(\sqrt{\delta^2 + (x-\mu)^2}/\alpha)^{\frac{1}{2}-\lambda}} \cdot e^{\beta(x-\mu)} \quad (2.1)$$

where  $K_\lambda$  is a modified Bessel function of the third kind with the index  $\lambda$ .

$$K_\lambda(\omega) = \frac{1}{2} \int_0^\infty \exp\left[-\frac{\omega}{2}(v^{-1} + v)\right] v^{\lambda-1} dv \quad (2.2)$$

In all cases,  $\mu$  is the location parameter and can take any real value,  $\delta$  is a scale parameter;  $\alpha$  and  $\beta$  determine the distribution shape and  $\lambda$  defines the subclasses of GH and is related to the tail flatness.

Many distributions are obtained as limiting distributions of the generalized hyperbolic distribution and by varying parameter  $\lambda$  to obtain subclasses for example, hyperbolic distribution and normal inverse Gaussian.

### 2.2.1 Hyperbolic distributions

When  $\lambda = 1$ , we obtain the subclass of hyperbolic distribution with probability density function (see Eberlein and Keller [1995],)

$$f_{hyp}(x; \alpha, \beta, \delta, \mu) = \frac{\sqrt{\alpha^2 - \beta^2}}{2\alpha\delta K_1(\delta\sqrt{\alpha^2 - \beta^2})} \exp\left[-\alpha\sqrt{\delta^2 + (x-\mu)^2} + \beta(x-\mu)\right]. \quad (2.3)$$

The mean and variance of hyperbolic function given respectively by the followings

$$E(X) = \mu + \frac{\beta\delta}{\sqrt{\alpha^2 - \beta^2}} \frac{K_2(\eta)}{K_1(\eta)} \quad (2.4)$$

$$Var(X) = \delta^2 \left( \frac{K_2(\eta)}{\eta K_1(\eta)} + \frac{\beta^2}{\alpha^2 - \beta^2} \left[ \frac{K_3(\eta)}{K_1(\eta)} - \left( \frac{K_2(\eta)}{K_1(\eta)} \right)^2 \right] \right) \quad (2.5)$$

where  $\eta = \delta\sqrt{\alpha^2 - \beta^2}$ . The first two parameters  $\alpha$  and  $\beta$  determine the shape of the distribution, while the other two  $\delta$  and  $\mu$  are scale and location parameters.

### 2.2.2 NIG distribution

A random variable  $X \sim NIG(\alpha, \beta, \delta, \mu)$  if ( see Barndorff-Nielsen [1998])

$$f_{NIG}(x; \alpha, \beta, \delta, \mu) = \frac{\alpha}{\pi} \exp\left(\delta \left[\sqrt{\alpha^2 - \beta^2} + \beta\zeta(x)\right]\right) \frac{K_1(\alpha\delta\sqrt{1 + \zeta(x)^2})}{\sqrt{1 + \zeta(x)^2}}$$

where  $\zeta(x) = (x - \mu)/\delta$  and  $K_1$  is the modified Bessel function of third kind, with the index 1.

$$K_1(\omega) = \frac{1}{2} \int_0^\infty \exp\left[-\frac{\omega}{2}(v^{-1} + v)\right] dv$$

Loglikelihood function for MLE of parameters

$$\begin{aligned} \mathcal{L}_{NIG}(x|\alpha, \beta, \delta, \mu) &= -n \ln\left(\frac{\pi}{\mu}\right) + n(\delta\sqrt{\alpha^2 - \beta^2}) - \frac{1}{2} \sum_{i=1}^n \log(1 + \zeta(x_i)^2) \\ &\quad + \beta\delta \sum_{i=1}^n \zeta(x_i) + \sum_{i=1}^n \log(K_1(\delta\alpha\sqrt{(1 + \zeta(x_i)^2)})) \end{aligned}$$

## 3 Modeling underlying process

Most of the notation and model framework in this paper, is a slight modification of the skewed and leptokurtic generalized GARCH framework proposed by Stentoft [2008] and references therein. In that study, the dynamics of log returns  $X_t$  are specified as

$$\begin{aligned} X_t &= m_t(\cdot; \theta_m) + \sigma_t \varepsilon_t \\ \sigma_t^2 &= g(\sigma_s^2, \varepsilon_s; -\infty < s \leq t - 1; \theta_h) \\ \varepsilon|\mathcal{F}_{t-1} &\sim D(0, 1; \theta_D) \end{aligned}$$

where  $\mathcal{F}_{t-1}$  is the information set containing all information up to and including time  $t - 1$ ,  $m_t(\cdot; \theta_m)$  denote the conditional mean, governed by a set of parameters  $\theta_m$ ,  $D(0, 1; \theta_D)$  denote a zero mean and a unit variance distribution function, allowed to depend on a set of parameters  $\theta_D$ .

Let  $(\Omega, \mathcal{F}, (\mathfrak{F}_t)_{t \in [0, T]}, \mathbb{P})$  be a stochastic basis describing the uncertainty of the economy. We refer to  $\mathbb{P}$  as the physical probability measure and  $\mathfrak{F}_t$  represent the information flow driven by Wiener process  $W = (W_t)_{t \in [0, T]}$  and Lévy proces  $L = (\mathfrak{L}_t)_{t \in [0, T]}$ . Let  $S_t$  be the price of a stock at time  $t$  adapted to the natural filtration  $\mathfrak{F}_t$ . Define daily log return as  $X_t = \log S_{t+d_t} - \log S_{t-1}$ ,  $t = 1, 2, \dots$  where  $d_t$  denote dividends at time  $t$  and  $X_t$  the continuously

**Theorem 3.1.** *We propose the following model for asset returns under  $\mathbb{P}$ . Let*

$$X_t = \log\left(\frac{S_t + d_t}{S_{t-1}} \middle| \mathcal{F}_{t-1}\right)$$

where  $d_t$  is one period dividends paid, then

$$\begin{aligned} X_t &= m_t(\cdot; \theta_m) + \sigma_t(W_t + \mathfrak{L}_t), \\ &= m_t(\cdot; \theta_m) + \sigma_t(\vartheta_t + \varpi\xi_t), \quad \xi \in GH \\ \sigma_t^2 &= g(\sigma_s^2, \vartheta_s; -\infty < s \leq t-1; \theta_\sigma), \\ &\quad \vartheta_t \sim i.i.d.(0, 1), \quad \xi_t \sim i.i.d.D(0, 1; \theta_D) \end{aligned}$$

where  $m_t(\cdot; \theta_m)$  denote the mean function,  $\sigma_t(\cdot; \theta_\sigma)$  denote the variance process and  $\theta = (\theta_m, \theta_\sigma, \theta_D)$ .

### 3.1 Examples from Benchmark models

Without loss of generality, for all the subsequent examples we assume  $d_t = 0 \forall t > 0$

**Example 3.1.** *Geometric Brownian motion*

Let  $dS_t = \mu S_t dt + \sigma S_t dW_t$  be the stochastic differential equation modeling the uncertainty of the underlying process, then

$$\begin{aligned} S_t &= S_0 \exp\left(\left[\mu - \frac{\sigma^2}{2}\right]t + \sigma\sqrt{t}Z\right), \quad Z \sim N(0, 1) \\ \Rightarrow X_t &= \log(S_t/S_{t-1}) = \left[\mu - \frac{\sigma^2}{2}\right] + \sigma Z \\ \text{thus} \quad X_t &= \left[\mu - \frac{\sigma^2}{2}\right] + \sigma(Z + 0\xi), \quad \theta_m = (\mu, \sigma^2), \\ X_t &= m_t(\cdot; \theta_m) + \sigma_t(Z + \varpi\xi), \quad m_t = \left[\mu - \frac{\sigma^2}{2}\right], \quad \sigma_t = \sigma, \quad Z \sim i.i.d.N(0, 1) \end{aligned}$$

**Example 3.2.** *Jump-diffusion model*

Mertons(1976) introduced and analyzed one of the first models with both jump and diffusion term for pricing of derivative securities. Merton jump-diffusion model can be specified through the SDE..

$$\frac{dS_t}{S_t} = \mu dt + \sigma dW_t + (J_t - 1)dN_t$$

where the jump sizes are  $J_t$  are identically distributed and mutually independent. He also assumed that the three processes  $(W_t)_{t \geq 0}$ ,  $(N_t)_{t \geq 0}$  and  $(J_t)_{t \geq 0}$  are independent. Let  $\sum_{j=1}^{N(t)} (Y_j - 1)$  where  $Y_1, Y_2, \dots$ , are random variables and  $N(t)$  is a counting process. The following expression solves the SDE.

$$S(t) = S(0) \exp\left[\left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t\right] \prod_{j=1}^{N(t)} Y_j$$

which implies

$$\begin{aligned} \log_e\left(\frac{S(t)}{S(0)}\right) &= \left(\mu - \frac{\sigma^2}{2}\right)t + \sigma W_t + \sum_{j=1}^{N_t} \log_e Y_j \\ &= \left(\mu - \frac{\sigma^2}{2}\right)t + \sigma \left(W_t + \frac{\varpi}{\sigma} \sum_{j=1}^{N_t} \log_e Y_j\right) \end{aligned}$$

Given any date  $t \geq 0$  and a holding period of length  $h > 0$ , the returns  $X_t(h)$  over the period  $[t, t+h]$  is a model given by

$$X_t(h) = \begin{cases} x, & \text{if } K = 0; \\ x + y_1 + \dots + y_k, & \text{if } K \geq 1. \end{cases} \quad (3.1)$$

where  $x \sim N(\alpha h, \sigma^2 h)$ ,  $\alpha = (\mu - (\sigma^2/2))$ ,  $y_1, \dots, y_k$  is an i.i.d. sequence with common distribution say  $G$  and  $K$  is Poisson with parameter  $\lambda h$ ,  $\lambda > 0$  for  $k = 0, 1, 2, \dots$ , we have  $\text{Prob}(K = k) = \exp(-\lambda h)(\lambda h)^k/k$ . If  $Y_j$  have the log normal distribution  $LN(a, b)$  then  $\log_e Y_j \sim N(a, b^2)$  and  $\sum \log Y_j \sim N(an, b^2 n)$  which is the same as  $\sum \log Y_j \sim an + b\sqrt{n}N(0, 1)$ . Thus

$$\begin{aligned} X_t &= \log_e \left( \frac{S_t}{S_{t-1}} \right) \\ &= \left( \mu - \frac{\sigma^2}{2} \right) + \sigma \left( Z_t + \frac{\eta}{\sigma} \sum_{j=1}^K \log_e Y_j \right) \\ &= m(\cdot, \theta_m) + \sigma_t(Z_t + \varpi \xi) \text{ where } \xi = \sum_{j=1}^K \log_e Y_j \end{aligned}$$

**Example 3.3.** *Duan(1995) GARCH model*

Discrete time economy, one period rate of return assumed to be conditionally lognormally distributed under  $\mathbb{P}$ . Let  $r$  be constant one period risk free rate of return,  $\lambda$  be constant unit risk premium then

$$\begin{aligned} X_t &= \log(S_t/S_{t-1}) \\ &= r + \lambda \sigma_t - \frac{\sigma_t^2}{2} + \sigma_t^2 Z, \text{ where} \\ \sigma_t^2 &= \omega + \sum_{j=1}^q \alpha_j \sigma_{t-j}^2 Z_{t-j} + \sum_{j=1}^p \beta_j \sigma_{t-j}^2, \quad Z_{t-j} \sim i.i.d.N(0, 1) \forall j \\ &= m_t(\cdot; \theta_m) + \sigma_t(\vartheta_t + \varpi \xi) \text{ where } \sigma_t \sim \text{GARCH}(p, q) \end{aligned}$$

**Example 3.4.** *It is well known from empirical studies that  $X_t$  can be represented as  $X_t = \mu_t + \varepsilon_t + \xi_t$  where  $\mu_t$  is a mean function and  $\varepsilon_t, \xi_t$  are the two components of the error term (see for instance Merton [1976], Heston and Nandi [2000], Maheu and McCurdy [2004], Mwaniki [2015]). Moreover, define a  $p^{\text{th}}$  order autoregressive process  $\{X_t, t \geq 0\}$  with  $\text{GARCH}(p, q)$  error as*

$$\begin{aligned} X_t &= \mu_t + \varepsilon_t + \xi_t, \text{ where} \\ \mu_t &= \sum_{r=1}^p \phi_r X_{t-r} + \mu, \quad t \in \mathbb{Z}^+ \\ \varepsilon_t + \xi_t &= \sigma_t(Z_t + \sigma \mathfrak{L}_t), \quad Z_t, \text{ and } \mathfrak{L}_t \sim i.i.d(0, 1), \quad Z_0 = 0, \mathfrak{L}_0 = 0 \\ \sigma_t^2 &= \text{GARCH}(p, q), \quad p, q \in \mathbb{Z}^+ \end{aligned} \quad (3.2)$$

where  $Z_t$  and  $\mathfrak{L}_t$  are identically and independently distributed random variables. A general time series model for log returns would be

$$\begin{aligned} X_t &= \mu_t + \sigma_t(Z_t + \sigma \mathfrak{L}_t), \quad Z_t \sim N(0, 1), \quad \mathfrak{L}_t \in GH \\ &= m(\cdot; \theta_m) + \sigma_t(Z_t + \varpi \mathfrak{L}_t) \end{aligned}$$

## 4 Empirical Analysis

We investigate the statistical properties under the objective measure  $\mathbb{P}$  model's ability to explain observed market share prices.

### 4.1 Data description and parameter estimates

We apply our framework to the stock indices sample from top GDP countries, S& P500 from New York Stock exchange, CAC40 Paris stock exchange. The financial time series data consist of S&P500 and CAC40 daily closing adjusted price from January 2, 2001 through December 31, 2014. Daily adjusted closing prices from were used to determine daily log returns  $X_t, t = 0, 1, 2, \dots$ . Let  $S_j$  be the price on day  $j, j = 0, 1, 2, \dots, n - 1$ . Sample increments of log returns is defined by  $X_j = \log S_j - \log S_{j-1}, j = 1, 2, \dots, n - 1$ . GARCH models are well known to be the best performing models to describe evolution of volatility, a satisfactory statistical fit is provided when the distribution of the filtered historical residuals is non-Gaussian. We fit  $AR(1) + GARCH(1, 1)$  model conditioned on normal distribution. All models parameters are estimated by numerical maximum likelihood routine and the significant parameters are reported in Table 1 and Table 2.

Table 1:  $AR(1) - GARCH(1, 1)$  model parameter estimates

S&P500	Estimate	Std. Error	t value	Pr(> t )	AIC	BIC	LL
$\bar{\mu}$	0.00054	0.00014	3.78729	0.00015	6.3902	6.3811	11251.68
$\phi$	-0.05695	0.01796	-3.17118	0.00152			
$\omega$	0.00000	0.00000	5.30935	0.00000			
$a$	0.09202	0.00925	9.94648	0.00000			
$b$	0.89493	0.00992	90.23790	0.00000			

Table 2:  $AR(1) - GARCH(1, 1)$  model parameter estimates

CAC40	Estimate	Std. Error	t value	Pr(> t )	AIC	BIC	LL
$\bar{\mu}$	0.00050	0.00018	2.81357	0.00490	5.92174	5.9131	10607.87
$\phi$	-0.05432	0.01765	-3.07695	0.00209			
$\omega$	0.00000	0.00000	4.44794	0.00001			
$a$	0.09129	0.00956	9.55340	0.00000			
$b$	0.89951	0.01012	88.84475	0.00000			

The standardized filtered residuals are known to be uncorrelated and weakly stationary in their first and second moments. To this end we make a simplifying assumption that the resulting sequence is independent and identically distributed from an unknown distribution. We fit Normal Inverse Gaussian and hyperbolic distribution to the residuals, estimated parameters of the two distribution are summarized in Table 3 and their corresponding kernel densities are compared in in Figure 1.

Table 3: NIG and Hyperbolic parameter estimates

	S&P500		CAC40	
	NIG	HYP	NIG	HYP
$\alpha$	1.60743	1.87176	1.99970	2.31156
$\beta$	-0.35232	-0.31274	-0.36798	-0.36449
$\delta$	1.50306	0.89654	1.89130	1.50307
$\mu$	0.29586	0.26133	0.30532	0.30219

## 4.2 Goodness of fit

Two sample test called Kolmogorov-Smirnov test (K-S test) was applied. Empirical CDF, an estimate of the underlying data is used to test the following hypothesis.

$$H_0 : F_n(x) = F(x) \text{ for all } x \text{ versus } H_1 : F_n(x) \neq F(x) \text{ for some } x$$

where  $F_n(x)$  is the empirical cumulative probability estimated as  $F_n(x_i) = i/n$  for the  $i$ th smallest data value.  $F(x)$  is the theoretical cumulative distribution function evaluated at  $x$ . The test statistic is given by

$$D_n = \sup_x |F_n(x) - F(x)|$$

If the null hypothesis is true, then the theoretical distribution fits very well. If  $D_n$  is sufficiently large; the null hypothesis can be rejected. For the two filtered residual we computed  $D_n$  statistics and got  $D = 0.0163$ ,  $pvalue = 1$ , for both indices (S&P500 and CAC40). This implies that the two sample sets seems to come from the same parent distribution. Attempts to test whether the unknown distributions came from either of the three proposed densities, was reported in Table 4. The graphical representation of Kernel densities in Figure 1, supports the claim that the two densities share the same parent distribution.

Table 4: Kolmogorov Smirnov distances

$K_{Dist}$	NIG	HYP	NORM
S&500	0.4872	0.4872	0.7136
CAC40	0.5077	0.5077	0.7091

Judged on the distance between the empirical and theoretical distributions of the residuals, the picture changes slightly. The Kolmogorov-Smirnov test allows us to reject all Gaussian models. As the Kolmogorov-Smirnov statistic measures the uniform distance between two distribution functions, it might be of interest to test the models ability to appropriately model leptokutic nature of log returns. If one suspects the data to be from a family of normal distribution, corresponding values for the Anderson - Darling distance which has its focus on the tails of the distribution will be valid. The following discrete version of the Anderson - Derling statistic AD, measuring the distance between the theoretical distribution function  $F$  and the empirical distribution  $\hat{F}$ :

$$AD(F, \hat{F}) = \sup_{x \in \mathbb{R}} \frac{|F(x) - \hat{F}(x)|}{\sqrt{F(x)(1 - F(x))}}$$






Figure 1: CAC40 filtered returns and S&P500 filtered returns density estimations

From our study, it was clear that the data in question was non normal. Models based on the NIG and hyperbolic distribution which was used to provide a flexible description for empirically observed conditioned leptokurtic residuals seemed to fit the data.

## 5 Conclusions

Since log-Levy models fall short of explaining the auto correlation of the absolute return, we have observed that that the log return dynamics can be modelled by assuming three components (the mean function, the volatility function and the GARCH filtered residuals). Identification of the probability distributions can be estimated by class of distributions which can capture skewness and kurtosis for example normal inverse Gaussian or hyperbolic distributions. It is widely recognized that the key to developing successful strategies for managing risk and pricing assets is to parsimoniously describe the stochastic process governing the asset dynamics. This paper proposes general framework assumed to improve modeling returns of financial time series data. To this end, the proposed framework combine two major stylized facts of returns: changing variance and presence of excess kurtosis in filtered returns. The framework can be studied further by allowing different distributions under the objective measure  $\mathbb{P}$ . Empirical investigation indicate that the all the required model parameters can be estimated form from historical data and the filtered residual are non-normal. Further refinement of the model is left for future studies.

## 6 Acknowledgements

Financial support from International Science Program (ISP-Sweden) in collaboration with Eastern African Universities Mathematical program EAUMP is greatly appreciated.

## References

- Barndorff-Nielsen, O. (1998). Process of Normal Inverse Gaussian type. *Finance Stochastics*, 2:41–68.
- Black, F. and Scholes, M. (1973). The pricing of Options and Corporate liabilities. *Journal of Political Economy*, 81:659–683.
- Bollerslev (1986). Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31:307–327.
- Bollerslev, T., Engle, R., and Nelson, D. (1994). *ARCH models. Handbook of econometrics*, volume 4.
- Carr, P., German, H., Madan, D., and Yor, M. (2002). The Fine Structure of Asset Returns: An Empirical Investigation. *Journal of Business*, 75:305–332.
- Carr, P. and Wu, L. (2004). Time-Changed Lévy processes and option pricing. *Journal of Financial Economics*, 71:113–141.
- Cont, R. (2001). Empirical properties of asset returns: stylized facts and statistical issues. *Quantitative Finance*, 1:223–226.
- Duan, J. (1995). The garch option pricing model. *Mathematical Finance*, 5:13–32.
- Eberlein, E. and Keller, U. (1995). Hyperbolic Distributions in Finance. *Bernaoulli*, 1:281–299.
- Engle, R. (1982). Autoregressive Conditional Heteroscedasticity With Estimates of Variance of United Kingdom Inflation. *Journal of Business and Economic Statistics*, 9:987–1008.
- Glosten, L., Jagannathan, R., and Runkle, D. (1993). The relationship between expected value and the volatility of the nominal excess returns on stocks. *Journal of Finance*, 48:1779–1801.
- Heston, S. and Nandi, S. (2000). A closed form garch option valuation model. *Review of Financial Studies*, 13:585–625.
- Maheu, J. and McCurdy, H. (2004). News arrival Jump dynamics and volatility components for individual stock returns. *The Journal of Mathematical Finance*, 59:755–793.
- Merton, R. (1976). Option pricing when underlying stocks are discontinuous. *Journal of Financial Economics*, 3:125–144.
- Mwaniki, I. (2015). Modeling returns and unconditional variance in risk neutral world for liquid and illiquid market. *Journal of Mathematical finance*, 5:15–25.

- Pagan, A. and Schwert, G. (1990). Alternative models for conditional stock volatility. *Journal of Econometrics*, 45:267–290.
- Rogers, L. C. G. and Zhang, L. (2011). An asset return model capturing stylized facts. *Mathematics and Financial Economics*, 5(2):101–119.
- Rydberg, T. (2000). Realistic statistical modelling of financial data. *International Statistical Review*, 68(3):233–258.
- Stentoft, L. (2008). American option pricing using garch models and normal inverse gaussian distribution. *Journal of Financial econometrics*, 6(4):540–582.
- Tsay, R. (2002). *Analysis of Financial time series*. Wiley, first edition.