

# The Variational Iteration Method for Analytic Treatment of Homogeneous and Inhomogeneous Partial Differential Equations

M. O. Olayiwola<sup>1,2</sup>

<sup>1</sup> Department of Mathematical and Physical Sciences  
Faculty of Basic and Applied Sciences  
College of Science, Engineering and Technology  
Osun State University, Osogbo, Nigeria  
<sup>2</sup> Information Management and Technology Centre  
Office of the Vice-Chancellor  
Osun State University, Osogbo, Nigeria

E-mail:- olayiwola.oyedunsi@uniosun.edu.ng  
Tel:-+2348028063936, +23408055218234, +2348131109234

**Abstract** In this paper, the author used the variational iteration method (VIM) to find the analytical solution to homogeneous and inhomogeneous partial differential equations. Few numerical examples were presented to show the effectiveness and efficiency of the method. It was observed that by carefully chosen a very good choice of initial guess leads to a solution in a closed form. The method is elegant and reliable.

**Keywords:** variational iteration method, initial value problems, partial differential equations.

## 1.0 Introduction

Partial differential equations; linear or nonlinear, homogenous or inhomogeneous has many applications to real life problems that arise in science, engineering and technology.

There are many numerical methods for the solution of different types of differential equations such as Adomian decomposition method [1-3], homotopy perturbation method [4], variational iteration method [5-9], modified variational iteration method 14-15]. Results by various researchers [1-15] have shown reliability, efficiency and applicability of these methods.

In this paper, a variational iteration method for the solution of homogeneous and inhomogeneous partial differential equations is presented. It is to be noted that the Lagrange multiplier reduces the iteration on integral operator and also minimizes the computational time. The method requires no transformation and or linearisation of any forms. Some numerical problems and results are presented to show the reliability of the method.

## 2.0 The Variational Iteration Method

The basic idea of the He's Variational Iteration Method (VIM) [5-9], can be explained by considering the following nonlinear partial differential equations

$$Lu + Nu = g(x) \dots\dots\dots(1)$$

Where L is the linear operator, N is the nonlinear operator and g(x) is the inhomogeneous term. According to the method, we can construct a correction functional as follows

The corresponding variational iteration method for solving (1) is given as

$$u_{n+1}(x) = u_n(x) + \int_0^x \lambda \left[ Lu_n(s) + N \bar{u}_n(s) - g(s) \right] ds, \dots \dots \dots (2)$$

where  $\lambda$  is a Lagrange multiplier which can be identified optimally by variational iteration method. The subscript  $n$  denote the  $n$ th approximation,  $\bar{u}_n$  is considered as a restricted variation i.e  $\delta \bar{u}_n = 0$ . The successive approximation  $u_{n+1}, n \geq 0$  of the solution  $u$  can be easily obtained by determine the Lagrange multiplier and the initial guess  $u_0$ , consequently, the solution is given by  $u = \lim_{n \rightarrow \infty} u_n$ .

### 3.0 Application and Numerical Results

In this section, six problems will be presented to illustrate the efficiency of the method.

#### Example 3.1

Consider the following inhomogeneous equation

$$u_x - u_y = 1 + 2x + 2y, \quad u(0, y) = y + y^2, u(x, 0) = 2x + 3x^2 \dots \dots \dots (3)$$

The correction functional is given by

$$u_{n+1}(x, y) = y + y^2 + \int_0^x \lambda(s) \left[ \frac{\partial u_n(s, y)}{\partial s} - \frac{\partial u_n(s, y)}{\partial y} - 1 - 2s - 2y \right] ds \dots \dots \dots (4)$$

Making the correction functional stationary to obtain  $\lambda(s) = -1$ , hence, the iterative formula becomes

$$u_{n+1}(x, y) = y + y^2 - \int_0^x \left[ \frac{\partial u_n}{\partial x} - \frac{\partial u_n}{\partial y} - 1 - 2x - 2y \right] ds \dots \dots \dots (5)$$

Consequently, following approximants are obtained

$$u_1(x, y) = y + y^2 + 4xy + x^2 + 2x \dots \dots \dots (6)$$

$$u_2(x, y) = y + y^2 + 4xy + x^2 + 2x \dots \dots \dots (7)$$

$$u_3(x, y) = y + y^2 + 4xy + x^2 + 2x \dots\dots\dots(8)$$

the closed form and exact solution is given as

$$u(x, y) = y + y^2 + 4xy + x^2 + 2x \dots\dots\dots(9)$$

**Example 3.2**

Consider the following inhomogeneous equation

$$u_x + u_y = 2xy^2 + 2x^2y, \quad u(0, y) = 0, u(x, 0) = 0 \dots\dots\dots(10)$$

The correction functional and iterative formula becomes:

$$u_{n+1}(x, y) = 0 - \int_0^x \lambda(s) \left[ \frac{\partial u_n(s, y)}{\partial s} + \frac{\partial u_n(s, y)}{\partial y} - 2sy^2 - 2s^2y \right] ds \dots\dots\dots(11)$$

then, following approximants are obtained

$$u_1(x, y) = x^2y^2 + \frac{2}{3}x^3y \dots\dots\dots(12)$$

$$u_2(x, y) = x^2y^2 + \frac{1}{6}x^4 \dots\dots\dots(13)$$

$$u_3(x, y) = x^2y^2 \dots\dots\dots(14)$$

$$u_4(x, y) = x^2y^2 \dots\dots\dots(15)$$

The exact solution is given by

$$\lim_{n \rightarrow \infty} u(x, y) = x^2y^2 \dots\dots\dots(16)$$

**Example 3.3**

Consider the following equation

$$u_x + u_y = 2u, \quad u(0, y) = e^y, u(x, 0) = e^x \dots\dots\dots(17)$$

The correction functional is given by

$$u_{n+1}(x, y) = e^y - \int_0^x \left[ \frac{\partial u_n(s, y)}{\partial s} + \frac{\partial u_n(s, y)}{\partial y} - 2u(s, y) \right] ds \dots\dots\dots(18)$$

from (18), following iterations are obtained

$$u_1(x, y) = e^y + xe^y \dots\dots\dots(19)$$

$$u_2(x, y) = e^y + xe^y + \frac{x^2}{2} e^y \dots\dots\dots(20)$$

$$u_3(x, y) = e^y + xe^y + \frac{x^2}{2} e^y + \frac{x^3}{6} e^y \dots\dots\dots(21)$$

.  
.  
.

The series solution is given by

$$\begin{aligned} u(x, y) &= e^y + xe^y + \frac{x^2}{2!} e^y + \frac{x^3}{3!} e^y + \frac{x^4}{4!} e^y + \frac{x^5}{5!} e^y + \frac{x^6}{6!} e^y + \dots \\ &= e^y \left( 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \dots \right) \dots\dots\dots(22) \end{aligned}$$

and the closed form solution is

$$u(x, y) = e^{x+y} \dots\dots\dots(23)$$

which is the exact solution.

**Example 3.4**

Consider the following equation

$$u_x + u_y + u_z = 3, \quad u(0, x, y) = y + z, u(x, 0, z) = x + z, u(x, y, 0) = x + y \dots\dots\dots(24)$$

The correction functional and formula is given by

$$u_{n+1}(x, y, z) = y + z - \int_0^x \left[ \frac{\partial u_n(s, y, z)}{\partial s} + \frac{\partial u_n(s, y, z)}{\partial y} + \frac{\partial u_n(s, y, z)}{\partial z} - 3 \right] ds \dots\dots\dots(25)$$

Consequently, following approximants are obtained

$$u_1(x, y) = x + y + z \dots\dots\dots(26)$$

$$u_2(x, y) = x + y + z \dots\dots\dots(27)$$

.  
.
   
.

$$u(x, y) = x + y + z \dots\dots\dots(28)$$

This is the exact solution to (24).

**Example 3.5**

Consider the following equation

$$u_x + u_y + u_z = 3u, \quad u(0, x, y) = e^{y+z}, u(x, 0, z) = e^{x+z}, u(x, y, 0) = e^{x+y} \dots\dots\dots(29)$$

The correction functional is given by

$$u_{n+1}(x, y, z) = e^{y+z} - \int_0^x \left[ \frac{\partial u_n(s, y, z)}{\partial s} + \frac{\partial u_n(s, y, z)}{\partial y} + \frac{\partial u_n(s, y, z)}{\partial z} - 3u(s, y, z) \right] ds \dots\dots\dots(30)$$

therefore,

$$u_1(x, y) = e^{y+z} + xe^{y+z} \dots\dots\dots(31)$$

$$u_2(x, y) = e^{y+z} + xe^{y+z} + \frac{x^2}{2!} e^{y+z} \dots\dots\dots(32)$$

$$u_3(x, y) = e^{y+z} + xe^{y+z} + \frac{x^2}{2!} e^{y+z} + \frac{x^3}{3!} e^{y+z} + \dots \dots\dots(33)$$

.  
.
   
.

The series solution is given by

$$u(x, y) = e^{y+z} \left( 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \dots \right) \dots\dots\dots(34)$$

and the closed form solution is

$$u(x, y) = e^{x+y+z} \dots\dots\dots(35)$$

**Example 3.6**

Consider the following equation

$$u_x - u_y = 0, \quad u(0, y) = \sin y, u(x,0) = \sin x \dots\dots\dots(36)$$

The correction functional is given by

$$u_{n+1}(x, y) = \sin y - \int_0^x \left[ \frac{\partial u_n(s, y)}{\partial s} + \frac{\partial u_n(s, y)}{\partial y} \right] ds \dots\dots\dots(37)$$

Consequently, following approximant are obtained

$$u_1(x, y) = \sin y + x \cos y \dots\dots\dots(38)$$

$$u_2(x, y) = \sin y + x \cos y - \frac{x^2}{2!} \sin y \dots\dots\dots(39)$$

$$u_3(x, y) = \sin y + x \cos y - \frac{x^2}{2!} \sin y - \frac{x^3}{3!} \cos y \dots\dots\dots (40)$$

$$u_4(x, y) = \sin y + x \cos y - \frac{x^2}{2!} \sin y - \frac{x^3}{3!} \cos y - \frac{x^4}{4!} \sin y \dots\dots\dots(41)$$

$$u_5(x, y) = \sin y + x \cos y - \frac{x^2}{2!} \sin y - \frac{x^3}{3!} \cos y - \frac{x^4}{4!} \sin y + \frac{x^5}{5!} \cos y \dots\dots\dots (42)$$

The series solution is given by

$$u(x, y) = \sin y \left( 1 - \frac{x^2}{2!} - \frac{x^4}{4!} + \dots \right) + \cos y \left( x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots \right) \dots\dots\dots(43)$$

and the closed form solution is

$$u(x, y) = \sin x \cos y + \sin y \cos x = \sin(x + y) \dots\dots\dots(44)$$

which is the exact solution.

**4.0 Conclusion**

The paper has successfully described and applied the variational iteration method to some partial differential equations of physical significance. The method provides the solutions in terms of rapidly convergent series. It is also clear and remarkable that approximate solutions are in good agreement with analytical solution. The VIM was used in a direct way without using linearization, perturbation or restrictive assumptions. The method is elegant and reliable.

## References

- [1]. G. Adomian (1978) : “Nonlinear dissipative wave equations” *Applied Mathematics Letters*, Vol. 11, pp 125-126.
- [2]. Hassan N .A Ismail, Kamal Raslam & Aziza A Abd Rabboh (2004) “Adomian Decomposition Method for Generalized Burger’s-Huxley and Burger’s-Fisher Equation.” *Applied Mathematics and Computation*. 159, 291-301.
- [3]. I.H Abdel-Halim Hassan (2008): “Comparison of differential transformation technique with Adomian decomposition method for linear and nonlinear initial value problems,” *Chaos, Soliton and Fractals*. Vol. 36, pp 53-465.
- [4]. Hossein Jafari etal (2008): Application of Homotopy Perturbation Method for Solving Gas Dynamic Equation. *Applied Mathematical Sciences*, Vol.2,2008. No 48,2393-2396.
- [5]. Inokuti M. (1978) “General use of the Lagrange multiplier in nonlinear mathematical physics in: S. Nemat Nasser (Ed), Variational method in the mechanics of solid,” *Pergamon Press*, 156-162.
- [6]. He. J.H (1998) “Approximate analytical solution for seepage flow with fractional derivatives in porous media” *Compt. Math. App. Mech. Eng.* 167. 57-68.
- [7].He. J.H (1998) “ Approximate solution of nonlinear differential equations with convolution product nonlinearities” *Comp. Math. Apply Mech. Eng.* 167: 69-73.
- [8].Ji-Huan He,(1999) “ Variational Iteration method: a kind of non-linear analytical technique: Some examples”, *Int. Journal of Non-linear mechanics*, 3494 ,699-708.
- [9].He. J.H (2000) ”Variational iteration method for autonomous ordinary differential system” *App. Maths and Computation*, 114, 115-123.
- [10]. Amit Goyal, Alka,Rama Gupta and C.Nagaraja Kumar (2011) “Solitary Wave Solutions for Burgers-Fisher type Equations with Variable Coefficients” *WASET* 60, 1742-1746.
- [11]. D.Kaya, S.M.El.Sayed,(2004) “A numerical simulation and explicit solutions of The generalized Berger-Fisher equation,” *Appl.Math. Comput.* 152, 403-413.
- [12]. H.N.A.Ismail,A.A.A.Rabboh (2004) “A restrictive Pade approximation for The solution of the generalized Fisher and Berger-Fisher equation” *Appl. Math. Comput.* 154, 203-210
- [13]. Javidi M (2006) “ Modified Pseudospectral Method for generalized Burger’s-Fisher Equation” *International Mathematical Forum*, 1, No. 32, 1555-1564
- [14] Olayiwola M O (2014) : “An Improved Algorithm for the Solution of Generalized Burger-Fishers Equations” *Applied Mathematics* Vol. 5, pp 1609-1614



[15] Olayiwola M O (2014)- Analytical Approximate to the Solution of Some Nonlinear Partial Differential Equations Journal of the Nigerian Association of Mathematical Physics Vol. 28(2), 2014,Pp 69-72.